


**SUPPLEMENTARY MATERIAL**


**corresponding to:**

**Identification and expression analysis of zebrafish  
*testis-specific gene 10 (tsga10)***


SHOHREH ASGHARI-GIVEHCHI<sup>1,2</sup> and MOHAMMAD HOSSEIN-MODARRESSI<sup>\*,3</sup>

<sup>1</sup>Department of Medical Genetics, School of Medicine, International Campus, Tehran University of Medical Sciences, Tehran, Iran, <sup>2</sup>Keenan Research Center for Biomedical Science, Li Ka Shing Knowledge Institute, St. Michael's Hospital, Toronto, Ontario, Canada and <sup>3</sup>Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

\*Address correspondence to: Mohammad Hossein-Modarressi. Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran. Tel: +98-218-895-3005. Fax: +98-218-895-3005. E-mail: modaresi@tums.ac.ir -  <https://orcid.org/0000-0003-2763-1964>



**Supplementary Fig. S1. Sense RNA control.** *One-cell stage (A), 12 hpf (B), 18 hpf (C).*



**Supplementary Fig. S2. Injected *tsga10* mRNA expression.** *The presence of injected mRNA was confirmed by RT-PCR. *tsga10* mRNA was expressed from one-cell stage to 48 hpf embryo after microinjection and gradually degraded after 48 hpf with a very weak expression at 72 hpf. Actin beta1 (*actb1*) was used as a control.*