

Gonadal defects in *Cited2*-mutant mice indicate a role for SF1 in both testis and ovary differentiation

ALEXANDER N. COMBES¹, CASSY M. SPILLER¹, VINCENT R. HARLEY², ANDREW H. SINCLAIR³,
SALLY L. DUNWOODIE^{4,5}, DAGMAR WILHELM¹ and PETER KOOPMAN^{*,1}

¹Division of Molecular Genetics and Development, Institute for Molecular Bioscience, The University of Queensland, Brisbane, QLD, 2Prince Henry's Institute of Medical Research, Melbourne, VIC, ³Murdoch Children's Research Institute, and Dept. of Paediatrics, University of Melbourne, Royal Children's Hospital, Melbourne, VIC, ⁴Developmental Biology Division, Victor Chang Cardiac Research Institute, Darlinghurst, Sydney, NSW and ⁵Faculty of Medicine, University of New South Wales, Kensington, Sydney, NSW, Australia

ABSTRACT Sex determination is regulated by a molecular antagonism between testis- and ovary-determining pathways in the supporting cell lineage of the gonadal primordia. Genes important for maintaining this lineage play critical roles in early gonadal development, but their influence on testis and ovary differentiation is unclear due to the severity of loss-of-function phenotypes. The transcription factor SF1 (Nr5a1/Ad4BP) is one such factor, required for establishing the supporting cell lineage, and for propagating the male pathway. In the gonad, *Sf1* expression is enhanced by the transcriptional co-factor Cited2. We have used the reduced levels of *Sf1* expression in *Cited2*^{-/-} mice as a hypomorphic model to gain insight into the sex-specific roles of SF1 function in gonadal development. In XY mutant mice, we found that testis development was delayed in *Cited2*^{-/-} gonads, and that testis structure was permanently disrupted. In XX *Cited2*^{-/-} gonads, ectopic cell migration was observed which correlated with a transient upregulation of *Fgf9*, and a delay in *Wnt4* then *Foxl2* expression. These data suggest a novel role for SF1 in promoting ovarian development in addition to its roles in testis differentiation.

KEY WORDS: *cited2*, *SF1*, *testis development*, *ovary development*, *antagonism*

Introduction

Sex determination is regulated by competing molecular pathways in the supporting cell lineage of the bipotential gonad (Capel, 2000). Some transcription factors expressed in this lineage play critical roles in the development of the genital ridge, yet their expression is maintained after sex determination. The extent to which these factors influence testis or ovary differentiation after the early gonad has formed remains unclear.

The genital ridges develop from the intermediate mesoderm on either side of the neural tube from 9 days post coitum (dpc) (Capel, 2000). Subsequently, genes promoting testis or ovarian development are expressed in the genital ridge to determine gonadal sex. The expression of wingless-related MMTV integration site 4 (*Wnt4*), R-spondin1 (*Rspo1*), and β-Catenin at this stage promotes ovary development while opposing testis development (Chassot *et al.*, 2008, Kim *et al.*, 2006, Liu *et al.*, 2009, Maatouk *et al.*, 2008, Tomizuka *et al.*, 2008, Vainio *et al.*, 1999). Conversely, expression of fibroblast growth factor 9 (*Fgf9*) promotes

testis development while repressing ovarian fate (Kim *et al.*, 2006). The expression of Sex determining region of the Y chromosome (*Sry*) in XY gonads tips the balance of these opposing signals towards testis development by upregulating *Sry*-related homeobox-containing gene 9 (*Sox9*), with assistance from steroidogenic factor 1/nuclear receptor subfamily 5, group A, member 1/adrenal 4-binding protein (SF1/Nr5a1/Ad4BP) (Sekido and Lovell-Badge, 2008). *Sox9* expression results in upregulation of paracrine signals FGF9 (Kim *et al.*, 2006) and prostaglandin D2 (Wilhelm *et al.*, 2007), which in turn, promote *Sox9* expression in undifferentiated somatic cells (Kim *et al.*, 2006; Moniot *et al.*, 2009; Wilhelm *et al.*, 2005), leading to repression of *Wnt4*. In an XX environment, in the absence of SRY, the WNT signal prevails to repress *Fgf9* expression and promote ovarian development (Kim *et al.*, 2006). Disruption or delay in expression of genes in either pathway can lead to partial or full development of the

Abbreviations used in this paper: dpc, days post coitum.

*Address correspondence to: Peter Koopman. Division of Molecular Genetics and Development, Institute for Molecular Bioscience, The University of Queensland, Brisbane, QLD 4072, Australia. Fax: +61-7-3346-2101. e-mail: p.koopman@imb.uq.edu.au - web: <http://www.imb.uq.edu.au/groups/koopman>

Accepted: 2 June 2009. Final author-corrected PDF published online: 24 July 2009.

opposing fate, resulting in sex reversal.

SF1 plays a number of important roles in gonadal development. In mice lacking *Sf1*, the genital ridges form initially, then progressively degenerate through apoptosis, and are entirely absent by 12.5 dpc (Luo *et al.*, 1994). SF1 is expressed in the genital ridge from 9.5 dpc, under the regulation of Wilms tumor suppressor protein 1 (WT1) (Wilhelm and Englert, 2002). Subsequently, SF1 acts as a cofactor to upregulate the male sex determining gene *Sox9* (Sekido and Lovell-Badge, 2008) and its target anti-Müllerian hormone (Arango *et al.*, 1999) to propagate the male pathway. SF1 is also expressed in steroidogenic cells in both sexes at later stages (Ikeda *et al.*, 2001). Currently it is unclear whether SF1 influences the transcriptional program of early ovarian development outside its role in steroidogenesis.

Regression of the gonads after sex determination in *Sf1* knockout mice prevents a full analysis of the roles of this gene in gonadal differentiation (Luo *et al.*, 1994; Sadovsky *et al.*, 1995). However, a conditional allele for this gene exists and has been used to study the cell-specific roles of *Sf1* (Jeyasuria *et al.*, 2004). Additionally, *Sf1* function is dose-sensitive, so *Sf1*-haploinsufficient mice have been used to gain insight into downstream target genes (Park *et al.*, 2005).

We reasoned that a hypomorphic mouse model might provide further insight into the functions of SF1 in gonadal differentiation. Mice lacking CBP/p300-interacting transactivator with Glu/Asp-rich carboxy-terminal domain 2 (*Cited2*) provide a suitable model as *Cited2* interacts with WT1 and enhances, but is not required for, gonadal expression of *Sf1* (Val *et al.*, 2007). *Cited2* is expressed briefly in the gonadal primordia at 10 dpc. However, after 10.5 dpc, expression in the gonad is minimal until it is expressed again at low levels at 13.5 dpc (Val *et al.*, 2007). *Sf1* expression is reduced in *Cited2*^{-/-} mice, resulting in a transcriptional delay of the male program of development. *Sry*, *Sox9*, and other male markers are expressed later and at lower levels than in wild type controls (Val *et al.*, 2007). However, these levels are corrected by 13.5 dpc, and gonads have been reported to develop normally from this point (Bamforth *et al.*, 2001; Val *et al.*, 2007). The brief window of *Cited2* expression early in gonadal development suggests that the delay in testis development in these mice (Val *et al.*, 2007) is caused by the loss of *Cited2*-mediated enhancement of *Sf1* levels, rather than a direct effect by loss of *Cited2*.

We used the low levels of *Sf1* in *Cited2*^{-/-} mice as a hypomorphic model to study the roles of SF1 in testis and ovary development. In agreement with published studies, testis development was delayed. However, we found that despite recovery at the

transcriptional level, testis cord organisation and morphology were more permanently disrupted. Further we uncovered a novel role for SF1 in promoting ovary development through enhancing expression of *Wnt4* and *Foxl2*. We found that the male pathway was not fully repressed in the early ovary. Consequently, *Fgf9* was transiently upregulated, and ectopic cell migration was induced before a resurgence of the female transcriptional program. These results reveal a novel role for SF1 in promoting ovarian development.

Results

Disrupted gonadal morphology in *Cited2*^{-/-} XY gonads

We first investigated gonadal morphology in the *Cited2* null mouse to determine whether the reported delay and recovery of gene expression driving testis development (Val *et al.*, 2007) also occurred at the cellular and morphological levels. Wild type (*Cited2*^{+/+}) and heterozygous knockout (*Cited2*^{+/-}) gonads were morphologically similar and were used interchangeably as controls when comparing testis structure.

At 12.5 dpc, *Cited2*^{+/+} testes had an established coelomic blood vessel (Fig. 1A, arrow) and distinct vasculature between clearly segmented testis cords (Fig. 1A, asterisks). In comparison, homozygous knockout (*Cited2*^{-/-}) XY gonads of the same age lacked testis cord organisation (Fig. 1D, brackets), and had a severely reduced, and discontinuous, coelomic blood vessel (Fig. 1D, arrow). By 13.5 dpc, when testis cords had become more regular in wild type gonads (Fig. 1B), knockout gonads of the same age were still severely disrupted (Fig. 1E). Large regions of the gonad remained disorganized (Fig. 1E, bracket), while in other




Fig. 1. Abnormal testis morphology in *Cited2*^{-/-} mice. (A-C) Representative images of gonadal morphology from 12.5–14.5 dpc *Cited2*^{+/+} and *Cited2*^{-/-} embryos. Asterisks indicate testis cords and an arrow indicates the coelomic vessel. (D-F) Abnormal morphology in *Cited2*^{-/-} gonads demonstrates that testis development is defective in embryos lacking this gene. Arrows indicate the coelomic vessel, brackets highlight regions where Sertoli and germ cells are not organised into testis cords, asterisks mark testis cords. A-F, Sertoli cells are marked by an anti-SOX9 antibody (green). A, B, D, E are confocal images of wholemount gonads; germ cells and vasculature are marked by an anti-PECAM-1 antibody (red). C and F are confocal images of paraffin sections; germ cells are marked by an anti-OCT4 antibody (red). $n \geq 3$ for each knockout stage analysed. Scale bar, 100 μ m.

areas, testis cords and the coelomic vessel had formed (Fig. 1E, asterisks and arrow). 13.5 dpc knockout testis morphology did not resemble 12.5 dpc wild type testis morphology, as would be expected if testis organisation was simply delayed. At 14.5 dpc, wild type gonads had increased in size and retained regular cord organisation (Fig. 1C), whereas knockout gonads were smaller and had a mixture of disorganised regions and areas where cord formation had occurred (Fig. 1F; $n \geq 3$ for each knockout stage analysed).

In summary, testis morphology was disrupted in XY gonads lacking *Cited2*, with no stage of cord development resembling wild type. Thus, although a delayed transcriptional program recovers to normal levels by 13.5 dpc (Val *et al.*, 2007), knockout testis morphology did not recover by 14.5 dpc, implicating *Cited2* and threshold levels of *Sf1* as a requirement for establishing normal testis morphology.

Ectopic cell migration in *Cited2*^{-/-} XX gonads


Because testis cord formation is critically dependent on endothelial cell migration from the mesonephros (Cool *et al.*, 2008; Combes *et al.*, 2009), we reasoned that a lack of cord organisation might indicate compromised cell migration into the testis. To test this hypothesis, we assessed the competency of 11.5 dpc *Cited2*^{+/+} and *Cited2*^{-/-} gonads and mesonephroi to induce and respond to cell migration cues respectively. We found that cells within *Cited2*^{-/-} mesonephroi were able to respond to migration cues from wild type testes (Fig. 2 A-C), and although *Cited2*^{-/-} testes were considerably smaller than their wild type or heterozygous counterparts, they were still able to induce cell migration from wild type mesonephroi (Fig. 2 D-F). Surprisingly, loss of one or two functional *Cited2* alleles from an XX gonad resulted in ectopic cell

migration (Fig. 2 G-I). To compare levels of cell migration in XX gonads of different genotypes we determined the number of GFP-positive units, as a proxy for the number of migrating cells, in the gonadal compartment of each XX gonad and plotted the individual, and averaged results (Fig. 2 J,K). Background levels of migration were observed in wild type XX gonads with an average of 6 ± 0.8 GFP-positive units detected per gonad ($n=5$). However, 24 ± 3.9 ($n=4$) units were detected in *Cited2*^{+/+} XX gonads, and this number doubled to 51 ± 8 ($n=5$) in *Cited2*^{-/-} XX gonads. Thus, loss of functional *Cited2* alleles, and the corresponding decrease in SF1, does not have an overt influence on cell migration in the XY gonad, but causes a dose-dependent de-repression of cell migration in the XX gonad.

Transient upregulation of the male pathway in *Cited2*^{-/-} XX gonads

Under normal conditions, mesonephric cell migration is repressed by WNT signalling in XX gonads as part of the program of ovarian development (Chassot *et al.*, 2008; Jeays-Ward *et al.*, 2003; Liu *et al.*, 2009; Tomizuka *et al.*, 2008). To determine whether the ectopic cell migration we observed correlated to a disruption or delay in the ovarian program, we assessed expression levels of *Sf1*, key initiators of the WNT signalling pathway (*Wnt4*, *Rspo1*), and the female marker gene forkhead box L2 (*Foxl2*) from 10.5 to 13.5 dpc in XX *Cited2*^{-/-} gonads (Fig. 3). In parallel, we assessed the male markers *Sox9* and *Fgf9* to test for ectopic activation of the male pathway.

As expected, *Sf1* expression was reduced in *Cited2*^{-/-} XX gonads at 10.5 and 11.5 dpc (Fig. 3A, B) in comparison to *Cited2*^{+/+} and *Cited2*^{+/+} littermates. However by 12.5 dpc, *Sf1* levels were similar between homozygous-null and wild type gonads (Fig. 3C),



gonads, compared to *Cited2*^{+/+} XX controls. Results for individual experiments are plotted as separate data points (+/ n=5, +/- n=4, -/ n=5). (K) Mean number of migrating cells from the data set used in J. Error bars represent the standard error of the mean. $p<0.007$ for the difference between *Cited2*^{+/+} and *Cited2*^{-/-}. Some images displayed in Figure 2 have been rotated and displayed on a black background for presentation purposes.

Fig. 2. *Cited2* represses ectopic migration in XX gonads but does not grossly influence migration in XY gonads. The influence of *Cited2* on mesonephric cell migration was determined by assaying *Cited2*^{+/+} (+/+), *Cited2*^{+/+} (+/-), and *Cited2*^{-/-} (-/-) mesonephroi (A-C), XY gonads (D-F), and XX gonads (G-I). Boxed text to the left of each row indicates the tissue in which the genotype was varied. The genotype of the varied tissue is indicated above each column. (A-C) Loss of *Cited2* from the mesonephros did not impair cell migration into an XY gonad $n>3$ for all. (D-F) Loss of *Cited2* from the XY gonad did not impair cell migration, although the size of the *Cited2*^{-/-} gonad was reduced $n>3$ for all. (G-I) Loss of *Cited2* from the XX gonad resulted in ectopic cell migration into *Cited2*^{+/+} gonads which was increased in *Cited2*^{-/-} gonads. An arrowhead in G indicates background levels of migration that occur under normal (*Cited2*^{+/+}) conditions. g, gonad; m, mesonephros. Scale bars, 50 μ m. (J) Quantitation of ectopic cell migration in multiple XX *Cited2*^{+/+} and *Cited2*^{-/-}

indicating that *Cited2* enhances initial expression of *Sf1*, but that continued *Sf1* expression is independent of *Cited2*.

Pro-ovary genes- *Wnt4*, *Rspo1*, and *Foxl2*-followed a similar trend to *Sf1*, being expressed at reduced levels at 10.5 dpc (Fig. 3 D,G,J). The reduction in *Wnt4* between wild type and null gonads was statistically significant ($p<0.031$), whereas the reduction in *Rspo1* and *Foxl2* was not. The reduced expression of these

genes persisted at 11.5 dpc with a significant reduction in levels of *Foxl2* (Fig. 3 E,H,K). By 12.5 dpc, levels of *Wnt4*, *Rspo1*, and *Foxl2* in *Cited2* null gonads were comparable to wild type levels, indicating a recovery of the female transcriptional program (Fig. 3 F,I,L). These data implicate *Sf1* in promoting the program of ovarian differentiation.

The decreased expression of *Wnt4*, *Rspo1*, and *Foxl2* corre-




Fig. 3. Expression profiling of *Sf1*, and key female and male markers in *Cited2* XX gonads. Quantitative real time RT-PCR for: (A-C) *Sf1*, (D-F) *Wnt4*, (G-I) *Rspo1*, (J-L) *Foxl2*, (M-O) *Sox9*, and (P-R) *Fgf9*, in *Cited2*^{+/+} (+/+), *Cited2*^{+/-} (+/-), and *Cited2*^{-/-} (-/-) XX gonads from 10.5 – 12.5 dpc. n = 3 for all samples and bars indicate the standard error of the mean.

lated with increased levels of *Sox9*, and a statistically significant increase of *Fgf9* in the same homozygous gonads (Fig. 3 M,N,P,Q). However the higher levels of expression were transient and reduced to those of wild type controls by 12.5 dpc (Fig. 3 O,R).

Overall, these data indicate that in the absence of *Cited2*, decreased *Sf1* levels cause a delay in the female program of development. This delay results in a transient upregulation of male markers and ectopic induction of cell migration, but is short-lived because the initial delay in the program of ovarian differentiation is corrected, or compensated for, by 12.5 dpc.

Discussion

In this study we used mice lacking *Cited2*, an established enhancer of *Sf1* expression, as a hypomorphic model to study the effect of reduced levels of *Sf1* on testis and ovarian development. We found that testis morphology was not only delayed, but severely disrupted, and did not recover by 14.5 dpc. Furthermore, we assayed cell migration and found that migration was not disrupted in XY gonads or mesonephroi lacking *Cited2*. However, ectopic cell migration was observed in the XX gonad in a dose-dependent response to loss of functional *Cited2* alleles. Correspondingly, the female pathway was initially delayed but recovered shortly after, implicating *Sf1* in the initiation of ovarian development. These data underscore the fragility of the molecular control of sex determination, in that reduced levels of *Sf1* in the male disrupt testis morphology, whereas in the female, promoters of the male pathway are not sufficiently suppressed. However, the recovery of both pathways at the transcriptional level demonstrates compensatory mechanisms that act to reinforce sexual differentiation in the face of disruptions.

A central assumption of our study is that *Cited2*^{-/-} mice are a suitable model in which to study gonadal development. Embryos lacking *Cited2* die between 13.5 and 17.5 dpc (Weninger *et al.*, 2005) with multiple developmental defects including cardiac malformations and exencephaly (Bamforth *et al.*, 2001, Barbera *et al.*, 2002, Yin *et al.*, 2002). Placental insufficiency, which impacts on embryo growth between 12.5 and 14.5 dpc, is the likely cause of death (Withington *et al.*, 2006). However, these defects do not appear to have a major influence on the early stages of gonadal development as testis morphology has previously been reported to be normal prior to death in embryos homozygous for a different *Cited2* knockout allele to the one used in this study (Bamforth *et al.*, 2001; Val *et al.*, 2007).

Instead, we view the delay in gonadal development observed in *Cited2*^{-/-} mice as a result of reduced levels of SF1. SF1 is known to promote the testis pathway through transcriptional regulation of key testis-determining genes (Arango *et al.*, 1999, Sekido and Lovell-Badge, 2008, Wilhelm and Englert, 2002). Genetic and biochemical evidence supports a role for *Cited2* together with *Wt1* in enhancing gonadal expression of *Sf1* (Buaas *et al.*, 2009, Val *et al.*, 2007). Further, we and others show that the delay and recovery of male (Val *et al.*, 2007), and female (this study) transcriptional programs correlate with a delay and recovery of *Sf1* levels. However, we cannot exclude the potential influence of low oxygen and nutrient levels caused by placental insufficiency in these mice (Withington *et al.*, 2006), nor the potential for *Cited2* to influence other genes important for testis or ovarian development.

We found that the disruptions in testis morphology were not due to a reduction in cell migration. What, then, is responsible for the defects in cord organisation seen in *Cited2*^{-/-} gonads? It is well established that threshold levels and correct timing of *Sry* and *Sox9* expression are required for initiation of the male pathway (Bullejos and Koopman, 2005, Hacker *et al.*, 1995). However, SOX9 protein was detected along the length of the *Cited2*^{-/-} gonad by 12.5 dpc (Fig. 1B), indicating significant activation of the male program throughout the testis at this stage. Furthermore, lower levels of *Sf1* and a delay in target gene expression in *Sf1*-haploinsufficient mice do not affect gross cord structure or function (Park *et al.*, 2005). It is possible that differences in SF1 levels, or background strains, between our *Cited2*^{-/-} model and the *Sf1*-haploinsufficient mice could result in a more severe delay in testis development and disrupted cord formation. Alternatively, the cord defects may be caused or amplified by deprivations resulting from placental insufficiency in these mice (Withington *et al.*, 2006), or by loss of an unknown *Cited2*-regulated factor required for cord organisation.

The disruption to development in *Cited2*^{-/-} ovaries suggests an additional and previously unsuspected role for SF1 in initiating ovary development through direct or indirect mechanisms. The induction of ectopic cell migration in the ovary indicates that one role of SF1 is to promote WNT signalling, which usually represses this event (Chassot *et al.*, 2008, Jeays-Ward *et al.*, 2003, Liu *et al.*, 2009, Tomizuka *et al.*, 2008). This link is supported by the reduced levels of *Wnt4*, and *Rspo1* in knockout gonads at 10.5 dpc. Levels of *Foxl2*, another major player in ovarian development (Ottolenghi *et al.*, 2007), are also reduced, suggesting that the entire program of ovarian differentiation is delayed. These results support a role for SF1 in promoting ovarian differentiation.

These data extend our understanding of SF1 as a factor required for early gonadal development by demonstrating clear consequences for testis and ovarian development in a hypomorphic *in vivo* model of *Sf1* expression. Our results affirm an important role for SF1 in ensuring normal testis differentiation and implicate this factor in regulating early ovarian development. Future experiments will be required to determine which components of the ovarian program are regulated by SF1.

Materials and Methods

Mouse strains

Embryos were collected from timed matings, with noon of the day on which the mating plug was observed designated 0.5 dpc. The AGFP transgenic mouse line constitutively expressing EGFP from an autosome (Hadjantonakis *et al.*, 1998) was a gift from Andras Nagy. The mouse line containing a null allele for *Cited2* on the C57Bl/6 background has been described previously (Barbera *et al.*, 2002). Embryo sex was determined by *Zfy* PCR (Hogan *et al.*, 1994).

Antibodies and immunofluorescence

The following antibodies were used: Rabbit anti-SOX9 antibody (Wilhelm *et al.*, 2005), Rat anti-platelet endothelial cell adhesion molecule-1/CD31 (PECAM-1, 1:200, BD Biosciences), Mouse anti-OCT4 (1:50, Santa Cruz Biotechnology). Secondary antibodies, goat anti-rat Alexa Fluor 594, goat anti-rabbit Alexa Fluor 488, and goat anti-mouse 594 were obtained from Invitrogen. Whole mount immunofluorescence (Combes *et al.*, 2009) and section immunofluorescence

(Wilhelm *et al.*, 2005) was performed as described.

Migration assay

Mice carrying the *Cited2* null allele were bred with AGFP mice to produce GFP-positive *Cited2*^{+/+} offspring. These mice were then used for timed pregnancies to obtain GFP-positive mesonephroi which were *Cited2*^{+/+} or *Cited2*^{-/-}. The mesonephric cell migration assay (Martineau *et al.*, 1997), adapted for use with GFP (Nishino *et al.*, 2000), was performed and quantified as described in (Combes *et al.*, 2009). Some images displayed in Fig. 2 have been rotated and displayed on a black background for presentation purposes.

RNA isolation and quantitative real time RT-PCR

Gonads (with mesonephroi at ≤ 11.5 dpc, without at > 11.5 dpc) were dissected from embryos in PBS and total RNA was immediately isolated using the Micro RNA kit (Qiagen) as per manufacturers' instructions, including the optional DNaseI/genomic DNA degradation step. cDNA was synthesised from 1 μ g of RNA by reverse transcription (Superscript III, Invitrogen) using random primers (Promega) according to manufacturers' instructions. The ABI Prism-7000 Sequence Detector System was used to analyse relative cDNA levels. Quantitative RT-PCR experiments were performed in triplicate and repeated on three independent biological samples each representing two pooled gonads. Results are represented as the average of the experimental value, with error estimated by standard error of the mean. Samples were analysed in 25 μ l reactions containing 1 μ l cDNA prepared as described above, SYBR Green PCR Master Mix (Applied Biosystems) and 1 μ l each of 3.75 μ M forward and reverse primers. Cycling conditions began with an initial 10 min step at 95°C followed by 40 cycles of 95°C for 15 sec and 60°C for 1 min in a two-step thermal cycle. Dissociation curves were analysed for each primer set and cDNA samples were normalised against β S rRNA using the $2^{-\Delta\Delta CT}$ method. Primer sequences (5' to 3' forward, 5' to 3' reverse) were:

18S	GATCCATTGGAGGGCAAGTCT, CCAAGATCCAACCTACGAGCTTTT;
Sf1	TTCGTCTGCTCAAGTCCCTCATC, CCTTACGAGGCTGTGGTTGTT;
Wnt4	CTGGACTCCCTCCCTGTCTTT, CATGCCCTTGTCACTGCAA;
Rspo1	CGACATGAACAAATGCATCA, CTCCTGACACTTGGTGCGAGA;
Foxl2	GCTACCCCGAGCCCCGAAGAC, GTGTTGTCCCGCCTCCCTTG;
Sox9	AGTACCCGCATCTGCACAAC, TACTTGTAAATGGGGTGGTCT;
Fgf9	CTATCCAGGGAACCAGGAAAGA, CTCGTTCATGCCGAGGTAGAG.

Acknowledgements

We thank Wendy Chua and Kylie Lopes Floro for technical assistance, and Josephine Bowles for critical comments on the manuscript. Confocal microscopy was performed at the ACRF/IMB Dynamic Imaging Centre for Cancer Biology, established with the support of the Australian Cancer Research Foundation. This work was supported by research grants from the Australian Research Council and the National Health and Medical Research Council of Australia.

References

- ARANGO, N.A., LOVELL-BADGE, R. and BEHRINGER, R.R. (1999). Targeted mutagenesis of the endogenous mouse Mis gene promoter: *in vivo* definition of genetic pathways of vertebrate sexual development. *Cell* 99: 409-419.
- BAMFORTH, S.D., BRAGANCA, J., ELORANTA, J.J., MURDOCH, J.N., MARQUES, F.I., KRANC, K.R., FARZA, H., HENDERSON, D.J., HURST, H.C. and BHATTACHARYA, S. (2001). Cardiac malformations, adrenal agenesis, neural crest defects and exencephaly in mice lacking *Cited2*, a new Tcfap2 co-activator. *Nat Genet* 29: 469-474.
- BARBERA, J.P., RODRIGUEZ, T.A., GREENE, N.D., WENINGER, W.J., SIMEONE, A., COPP, A.J., BEDDINGTON, R.S. and DUNWOODIE, S. (2002). Folic acid prevents exencephaly in *Cited2* deficient mice. *Hum Mol Genet* 11: 283-293.
- BUAAS, F.W., VAL, P. and SWAIN, A. (2009). The transcription co-factor CITED2 functions during sex determination and early gonad development. *Hum Mol Genet*.
- BULLEJOS, M. and KOOPMAN, P. (2005). Delayed Sry and Sox9 expression in developing mouse gonads underlies B6-Y(DOM) sex reversal. *Dev Biol* 278: 473-481.
- CAPEL, B. (2000). The battle of the sexes. *Mech Dev* 92: 89-103.
- CHASSOT, A.A., RANC, F., GREGOIRE, E.P., ROEPERS-GAJADIEN, H.L., TAKETO, M.M., CAMERINO, G., DE ROOIJ, D.G., SCHEDL, A. and CHABOISSIER, M.C. (2008). Activation of beta-catenin signaling by Rsp01 controls differentiation of the mammalian ovary. *Hum Mol Genet* 17: 1264-1277.
- COMBES, A.N., WILHELM, D., DAVIDSON, T., DEJANA, E., HARLEY, V., SINCLAIR, A. and KOOPMAN, P. (2009). Endothelial cell migration directs testis cord formation. *Dev Biol* 326: 112-120.
- COOL, J., CARMONA, F.D., SZUCSIK, J.C., CAPEL, B. (2008). Peritubular myoid cells are not the migrating population required for testis cord formation in the XY gonad. *Sex Dev* 2: 128-33.
- HACKER, A., CAPEL, B., GOODFELLOW, P. and LOVELL-BADGE, R. (1995). Expression of Sry, the mouse sex determining gene. *Development* 121: 1603-1614.
- HADJANTONAKIS, A.K., GERTSENSTEIN, M., IKAWA, M., OKABE, M. and NAGY, A. (1998). Generating green fluorescent mice by germline transmission of green fluorescent ES cells. *Mech Dev* 76: 79-90.
- HOGAN, B., BEDDINGTON, R., COSTANTINI, F. and LACY, E. (1994). *Manipulating the mouse embryo: A laboratory manual*. Cold Spring Harbor Laboratory Press. pp 497., Cold Spring Harbor, New York.
- IKEDA, Y., TAKEDA, Y., SHIKAYAMA, T., MUKAI, T., HISANO, S. and MOROHASHI, K.I. (2001). Comparative localization of Dax-1 and Ad4BP/SF-1 during development of the hypothalamic-pituitary-gonadal axis suggests their closely related and distinct functions. *Dev Dyn* 220: 363-376.
- JEAYS-WARD, K., HOYLE, C., BRENNAN, J., DANDONNEAU, M., ALLDUS, G., CAPEL, B. and SWAIN, A. (2003). Endothelial and steroidogenic cell migration are regulated by WNT4 in the developing mammalian gonad. *Development* 130: 3663-3670.
- JEYASURIA, P., IKEDA, Y., JAMIN, S.P., ZHAO, L., DE ROOIJ, D.G., THEMHEN, A.P., BEHRINGER, R.R. and PARKER, K.L. (2004). Cell-specific knockout of steroidogenic factor 1 reveals its essential roles in gonadal function. *Mol Endocrinol* 18: 1610-1619.
- KIM, Y., KOBAYASHI, A., SEKIDO, R., DINAPOLI, L., BRENNAN, J., CHABOISSIER, M.C., POULAT, F., BEHRINGER, R.R., LOVELL-BADGE, R. and CAPEL, B. (2006). Fgf9 and Wnt4 act as antagonistic signals to regulate mammalian sex determination. *PLoS Biol* 4: e187.
- LIU, C.F., BINGHAM, N., PARKER, K. and YAO, H.H. (2009). Sex-specific roles of beta-catenin in mouse gonadal development. *Hum Mol Genet* 18: 405-417.
- LUO, X., IKEDA, Y. and PARKER, K.L. (1994). A cell-specific nuclear receptor is essential for adrenal and gonadal development and sexual differentiation. *Cell* 77: 481-490.
- MAATOUK, D.M., DINAPOLI, L., ALVERS, A., PARKER, K.L., TAKETO, M.M. and CAPEL, B. (2008). Stabilization of beta-catenin in XY gonads causes male-to-female sex-reversal. *Hum Mol Genet* 17: 2949-2955.
- MARTINEAU, J., NORDQVIST, K., TILMANN, C., LOVELL-BADGE, R. and CAPEL, B. (1997). Male-specific cell migration into the developing gonad. *Curr Biol* 7: 958-968.
- MONIOT, B., DECLOSMENIL, F., BARRIONUEVO, F., SCHERER, G., ARITAKE, K., MALKI, S., MARZI, L., COHEN-SOLAL, A., GEORG, I., KLATTIG, J. *et al.* (2009). The PGD2 pathway, independently of FGF9, amplifies SOX9 activity in Sertoli cells during male sexual differentiation. *Development* 136: 1813-1821.
- NISHINO, K., KATO, M., YOKOUCHI, K., YAMANOUCHI, K., NAITO, K. and TOJO, H. (2000). Establishment of fetal gonad/mesonephros coculture system using EGFP transgenic mice. *J Exp Zool* 286: 320-327.
- OTTOLENGHI, C., PELOSI, E., TRAN, J., COLOMBINO, M., DOUGLASS, E., NEDOREZOV, T., CAO, A., FORABOSCO, A. and SCHLESSINGER, D.

- (2007). Loss of Wnt4 and Foxl2 leads to female-to-male sex reversal extending to germ cells. *Hum Mol Genet* 16: 2795-2804.
- PARK, S.Y., MEEKS, J.J., RAVEROT, G., PFAFF, L.E., WEISS, J., HAMMER, G.D. and JAMESON, J.L. (2005). Nuclear receptors SF1 and Dax1 function cooperatively to mediate somatic cell differentiation during testis development. *Development* 132: 2415-2423.
- SADOVSKY, Y., CRAWFORD, P.A., WOODSON, K.G., POLISH, J.A., CLEMENTS, M.A., TOURTELLOTTE, L.M., SIMBURGER, K. and MILBRANDT, J. (1995). Mice deficient in the orphan receptor steroidogenic factor 1 lack adrenal glands and gonads but express P450 side-chain-cleavage enzyme in the placenta and have normal embryonic serum levels of corticosteroids. *Proc Natl Acad Sci USA* 92: 10939-10943.
- SEKIDO, R. and LOVELL-BADGE, R. (2008). Sex determination involves synergistic action of SRY and SF1 on a specific Sox9 enhancer. *Nature* 453: 930-934.
- TOMIZUKA, K., HORIKOSHI, K., KITADA, R., SUGAWARA, Y., IBA, Y., KOJIMA, A., YOSHITOME, A., YAMAWAKI, K., AMAGAI, M., INOUE, A. et al. (2008). R-spondin1 plays an essential role in ovarian development through positively regulating Wnt-4 signaling. *Hum Mol Genet* 17: 1278-1291.
- VAINIO, S., HEIKKILA, M., KISPERT, A., CHIN, N. and MCMAHON, A.P. (1999). Female development in mammals is regulated by Wnt-4 signalling. *Nature* 397: 405-409.
- VAL, P., MARTINEZ-BARBERA, J.P. and SWAIN, A. (2007). Adrenal development is initiated by Cited2 and Wt1 through modulation of Sf-1 dosage. *Development* 134: 2349-2358.
- WENINGER, W.J., LOPES FLORO, K., BENNETT, M.B., WITHERINGTON, S.L., PREIS, J.I., BARBERA, J.P., MOHUN, T.J. and DUNWOODIE, S.L. (2005). Cited2 is required both for heart morphogenesis and establishment of the left-right axis in mouse development. *Development* 132: 1337-1348.
- WILHELM, D. and ENGLERT, C. (2002). The Wilms tumor suppressor WT1 regulates early gonad development by activation of SF1. *Genes Dev* 16: 1839-1851.
- WILHELM, D., HIRAMATSU, R., MIZUSAKI, H., WIDJAJA, L., COMBES, A.N., KANAI, Y. and KOOPMAN, P. (2007). SOX9 regulates prostaglandin D synthase gene transcription *in vivo* to ensure testis development. *J Biol Chem* 282: 10553-10560.
- WILHELM, D., MARTINSON, F., BRADFORD, S., WILSON, M.J., COMBES, A.N., BEVERDAM, A., BOWLES, J., MIZUSAKI, H. and KOOPMAN, P. (2005). Sertoli cell differentiation is induced both cell-autonomously and through prostaglandin signaling during mammalian sex determination. *Dev Biol* 287: 111-124.
- WITHERINGTON, S.L., SCOTT, A.N., SAUNDERS, D.N., LOPES FLORO, K., PREIS, J.I., MICHALICEK, J., MACLEAN, K., SPARROW, D.B., BARBERA, J.P. and DUNWOODIE, S.L. (2006). Loss of Cited2 affects trophoblast formation and vascularization of the mouse placenta. *Dev Biol* 294: 67-82.
- YIN, Z., HAYNIE, J., YANG, X., HAN, B., KIATCHOOSAKUN, S., RESTIVO, J., YUAN, S., PRABHAKAR, N.R., HERRUP, K., CONLON, R.A. et al. (2002). The essential role of Cited2, a negative regulator for HIF-1alpha, in heart development and neurulation. *Proc Natl Acad Sci USA* 99: 10488-10493.

Further Related Reading, published previously in the *Int. J. Dev. Biol.*

See Special Issue ***Pattern Formation*** edited by Michael K. Richardson and Cheng-Ming Chuong at:

<http://www.ijdb.ehu.es/web/contents.php?vol=53&issue=5-6>

The spatio-temporal pattern of testis organogenesis in mammals - insights from the mole

Francisco D. Carmona, Darío G. Lupiáñez, José-Ezequiel Martín, Miguel Burgos, Rafael Jiménez and Federico Zurita

Int. J. Dev. Biol. (2009) 53: 1035-1044

Foetal germ cells: striking the balance between pluripotency and differentiation

Patrick Western

Int. J. Dev. Biol. (2009) 53: 393-409

Aard is specifically up-regulated in Sertoli cells during mouse testis differentiation

Terje Svøgen, Annemiek Beverdam, Pali Verma, Dagmar Wilhelm and Peter Koopman
Int. J. Dev. Biol. (2007) 51: 255-258

Effects of FGF9 on embryonic Sertoli cell proliferation and testicular cord formation in the mouse

Louise Willerton, Robert A. Smith, David Russell and Sarah Mackay
Int. J. Dev. Biol. (2004) 48: 637-643

Developmental regulation of expression of Ran/M1 and Ran/M2 isoforms of Ran-GTPase in mouse testis


Pedro P López-Casas, Luis A López-Fernández, Mario Párraga, Dora B Krimer and Jesús del Mazo
Int. J. Dev. Biol. (2003) 47: 307-310

Differentiation of mouse primordial germ cells into female or male germ cells

N Nakatsuji and S Chuma
Int. J. Dev. Biol. (2001) 45: 541-548

Wilms' tumor suppressor gene (WT1) as a target gene of SRY function in a mouse ES cell line transfected with SRY.

Y Toyooka, S S Tanaka, O Hirota, S Tanaka, N Takagi, K Yamanouchi, H Tojo and C Tachi
Int. J. Dev. Biol. (1998) 42: 1143-1151



5 yr ISI Impact Factor (2008) = 3.271

