

The subterranean catfish *Phreatobius cisternarum* provides insights into visual adaptations to the phreatic environment

LOUISE N. PEREZ, BERTHA R. MARILUZ, JAMILY LOREN, AMY LIU,
MARCOS P. SOUSA, RODRIGO A. P. MARTINS JOHN S. TAYLOR and PATRICIA N. SCHNEIDER

ACCEPTED MANUSCRIPT

The International Journal of Developmental Biology publishes an “Accepted” manuscript format as a free service to authors in order to expedite the dissemination of scientific findings to the research community as soon as possible after acceptance following peer review and corresponding modification (where appropriate).

An “Accepted” manuscript is published online prior to copyediting, formatting for publication and author proofing, but is nonetheless, fully citable through its Digital Object Identifier (doi®). Nevertheless, this “Accepted” version is NOT the final version of the manuscript.

When the final version of this paper is published within a definitive issue of the journal with copyediting, full pagination, etc. the new final version will be accessible through the same doi and this "Accepted" version of the paper will disappear.

The subterranean catfish *Phreatobius cisternarum* provides insights into visual adaptations to the phreatic environment

Louise N. Perez^{1a}, Bertha R. Mariluz^{1a}, Jamily Lorena², Amy Liu³, Marcos P. Sousa⁴, Rodrigo A. P. Martins⁵ John S. Taylor³ and Patricia N. Schneider^{1*}

¹Instituto de Ciências Biológicas, Universidade Federal do Pará (UFPA), Belém, Pará, Brazil.

²Instituto Tecnológico Vale, Pará, Brazil

³Department of Biology, University of Victoria, Victoria, BC, Canada.

⁴Coordenação de Zoologia, Museu Paraense Emílio Goeldi (MPEG), Belém, Pará, Brazil.

⁵Programa de Biologia Celular e do Desenvolvimento, Instituto de Ciências Biomédicas, Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil.

Running title: Visual adaptations of the subterranean catfish *Phreatobius cisternarum*

Keywords: opsins; blind catfish; neuromasts; sensory system

^aAuthors contributed equally

***Correspondence:**

Patricia N. Schneider, Centro de Genômica e Biologia de Sistemas. Instituto de Ciências Biológicas, Universidade Federal do Pará, Rua Augusto Corrêa #01, Belém, PA, 66075-900, Brasil. E-mail: patricia.schneider@northwestern.edu, schneider@ufpa.br

Abstract

Vertebrate eyes share the same general organization, though species have evolved morphological and functional adaptations to diverse environments. Cave-adapted animals are characterized by a variety of features including eye reduction, loss of body pigmentation, and enhanced non-visual sensory systems. Species that live in perpetual darkness have also evolved sensory mechanisms that are independent of light stimuli. The subterranean catfish *Phreatobius cisternarum* lives in the Amazonian phreatic zone and displays a diversity of morphological features that are similar to those observed in cavefish and appear to be adaptations to life in the dark. Here we combine histological and transcriptome analyses to characterize sensory adaptations of *P. cisternarum* to the subterranean environment. Histological analysis showed that the vestigial eyes of *P. cisternarum* contain a rudimentary lens. Transcriptome analysis revealed a repertoire of eleven visual and non-visual opsins and the expression of 36 genes involved in lens development and maintenance. In contrast to other cavefish species, such as *Astyanax mexicanus*, *Phreatichthys andruzzii*, *Sinocyclocheilus anophthalmus* and *Sinocyclocheilus microphthalmus*, DASPEI neuromast staining patterns did not show an increase in the number of sensory hair cells. Our work reveals unique adaptations in the visual system of *P. cisternarum* to underground habitats and helps to shed light into troglomorphic attributes of subterranean animals.

Introduction

Hypogean species are characterized by the reduction or loss of eyes and pigments, accompanied by enhancement of other sensory structures (Jeffery, 2009; Mohun *et al.*, 2010; Partha *et al.*, 2017; Protas and Jeffery, 2012). In cave-adapted species, eye degeneration appears to be controlled by at least three different genetic programs. In the blind cavefish *Astyanax mexicanus*, degeneration is triggered by, among other mechanisms, *shh* overexpression and subsequent induction of lens cells apoptosis (Yamamoto *et al.*, 2004). Recent studies have shown that the retinal pigment epithelium (RPE) is critical for eye maintenance (Ma *et al.*, 2020). In the blind cave-dwelling fish *Phreatichthys andruzzii*, eye degeneration is controlled by the

apoptosis of retinal neurons, whereas in the eyeless golden-line fish *Sinocyclocheilus anophthalmus*, transcriptome analysis showed that retinal degeneration is associated with downregulation of transcription factors controlling opsin expression.

Surface fishes can detect moving and vibrating water objects using the mechanosensory organs, the neuromasts. These units are distributed alongside the lateral line and is also observed in the head, where it is called cephalic lateral line (Suppl. Fig 2) (Soares and Niemiller, 2018). Notably, to offset the loss of vision, other sensory functions have been enhanced in cavefish species, including those involving neuromasts, barbels, and taste buds (Meng *et al.*, 2013; Stemmer *et al.*, 2015).

The genus *Phreatobius* (Siluriformes) includes three species: *P. sanguijuella*, *P. dracunculus* and *P. cisternarum* (Fernandez *et al.*, 2007; Muriel-Cunha and Pinna, 2005; Shibatta *et al.*, 2007; Trajano, 2007). These species occur in the Amazon and Paraguay river basins, however, reports regarding their biology, reproductive cycle, ecology, and physiology remain scarce (Fernandez *et al.*, 2007; Muriel-Cunha and Pinna, 2005; Ohara *et al.*, 2016). *Phreatobius cisternarum* inhabits the superficial phreatic layer, which corresponds to the upper saturated stratum of soil found in hand-dug wells. This species is characterized by small eyes and reduction in both body size and pigmentation (**Fig. 1 A,B, Suppl. Fig. 1 A,B**). Besides the six protruding barbels characteristic of catfishes, *P. cisternarum* has a paddle-shaped caudal region formed by a caudal-fin extended dorsally and ventrally by numerous large procurrent rays, ventrally confluent with the anal fin, a strongly prognathous lower jaw, hypertrophied jaw muscles, and a bright red coloration (Shibatta *et al.*, 2007).

Opsins are a diverse group of G-protein-coupled receptors (GPCRs) broadly categorized as 'visual' or 'non-visual'. The best known opsins are visual opsins found in the rod and cone photoreceptor cells of the retina. These opsins detect light when their ligand, a chromophore called 11-*cis* retinal, undergoes photoisomerization. Fish are notable in having large visual and non-visual opsin repertoires compared to other vertebrates (Beaudry *et al.*, 2017; Lin *et al.*, 2017; Rennison *et al.*, 2012). For example, the four-eyed fish, *Anableps anableps*, and the distantly related zebrafish (*Danio rerio*), have ten visual opsins (Davies

et al., 2015; Owens *et al.*, 2009). Zebrafish also have 32 non-visual opsins (Davies et al., 2015). It is clear that such a large repertoire is not essential for vision or circadian rhythm regulation, as other animals have much smaller repertoires (Scholtyßek and Kelber, 2017). By surveying the transcriptome of *P. cisternarum*, we characterized the molecular as well as the morphological response of the visual system to life in perpetual darkness. Furthermore, we used a set of histological analysis, immunofluorescence, and neuromast staining approaches to study adaptations of the sensory system in this subterranean species.

Material and Methods

Specimen collection

Fifteen *P. cisternarum* adults, ranging from 30 mm to 58 mm, were collected from artificial wells in Benevides (Pará, Brazil) and maintained in dark with daily feeding. In addition to the light protected tanks, the light in the room was controlled to keep the fish in the perpetual dark. During feeding, tank protection was removed and lights were kept off. Three specimens of *Poecilia reticulata* were obtained in the pet trade and were maintained in individual tanks in a recirculating freshwater system at 24 to 28 °C with aeration at the Genomics and Systems Biology Center. Experiments and animal care were performed following animal care guidelines approved by the Animal Care Committee at the Universidade Federal do Para (protocol no. 037-2015).

Library preparation and Illumina sequencing

One specimen of *P. cisternarum* was anesthetized in 0.1% tricaine solution (Sigma-Aldrich). Total RNA was extracted from its head using TRIzol® (Life Technologies), following manufacturer's protocol. Paired-end 150bp libraries were generated with NEXTSeq Mid Output, according to standard protocol (Illumina) and sequenced on Illumina NEXTSeq 500 platform (NCBI Sequence Read Archive project: PRJNA491408 and run: SRR7878036).

De novo transcriptome assembly and gene annotation

Over 190 million raw sequence reads were obtained. Trimmomatic was used with default parameters to remove adapters (Bolger *et al.*, 2014). *De novo*

transcriptome assembly was performed using standard parameters in Trinity (Grabherr et al., 2011). The completeness of the assembled transcripts was assessed through the Benchmarking Universal Single-Copy Orthologs tool (BUSCO) (Seppey et al., 2019). Mapping was performed by means of Bowtie (Langmead and Salzberg, 2012) using the constructed transcriptome as a reference. StringTie (Pertea et al., 2015) was used to estimate the abundance of transcripts in Transcripts Per Million (TPM). The identification of lens and RPE genes was performed via tBLASTn search tool, and each contig of interest was searched with manual queries of the consensus sequences from Uniprot (Priyam et al., 2019), and ExPASy Translate predicted Open Reading Frames (ORFs) (Gasteiger et al., 2005). BLASTp (NCBI) was performed to confirm protein homology.

Characterization of visual and non-visual opsins

Forty-two zebrafish (*Danio rerio*) opsins (Davies et al., 2015) were used as query sequences in a tBLASTx survey of the *P. cisternarum* transcriptome (E-value of 10^{-6}). *Danio rerio* opsins were also used to survey the transcriptome of another species in the order Siluriformes, the channel catfish (*Ictalurus punctatus* (accession GCF_001660625.1_lpCoco_1.2)). Opsins from *P. cisternarum*, *I. punctatus* and the 42 *D. rerio* opsins were aligned using ClustalW in BioEdit (Hall, 1999; Thompson et al., 1994). Introns in the *P. cisternarum* opsin transcripts that were obvious in this alignment were removed; precise intron-exon boundaries could be identified by comparing the *P. cisternarum* sequences to *D. rerio* orthologs on the NCBI Graphical Sequence Viewer. The alignment was then truncated at the 5' and 3' ends because the degree of sequence divergence in these regions made alignments (even when translated into amino acids sequences) unreliable. Maximum Parsimony (MP) phylogenetic trees were reconstructed using MegaX version 10.1.8 (Stecher et al., 2020) and confidence in the topology was evaluated by Bootstrap re-analyses.

DASPEI staining

The fluorescent dye 2-[4-(di-methylamino)styryl]-N-ethylpyridinium iodide (DASPEI; Invitrogen) was used as a vital dye to stain hair cells within

neuromasts. Three *P. cisternarum* and three *P. reticulata* specimens were incubated in an embryo medium containing 0.005% DASPEI for 15 min, anesthetized in 0.1% tricaine solution (Sigma-Aldrich) for 5 minutes, and rinsed once in fresh embryo medium according to a previously established protocol (Yoshizawa *et al.*, 2010). The specimens were analyzed and photographed with a NIKON-Eclipse 80i fluorescence microscope using the NIS-Elements imaging software.

Histological analysis

Twelve specimens of *P. cisternarum* were anesthetized in 0.1% tricaine solution (Sigma-Aldrich). The heads were collected and flash-frozen in Tissue-Tek embedding medium (Sakura). Cryosections (20 μm) were obtained on a Leica CM1850 UV cryostat (Leica Biosystems), captured on Color Frost Plus microscope slides (Thermo Fisher Scientific), fixed in 3% paraformaldehyde, and stored at -80°C for further use. Slides were stained with hematoxylin and eosin following standard protocol (Kiernan, 2008).

Immunostaining

Slides were incubated with gamma-crystallin antibody (a gift from Dr. Martins) overnight at 4°C following previously established protocol (Cavalheiro *et al.*, 2014). Immunofluorescence reaction was performed with an anti-rabbit secondary antibody (1:200, Sigma-Aldrich, cat# F6005), incubated for 2 hours at room temperature. Fluorescence nuclear counterstaining was performed with DAPI (Sigma-Aldrich, cat# F6057). Images were captured with a NIKON-Eclipse 80i fluorescence microscope and NIS-Elements imaging software.

Results and Discussion

Opsin repertoire in P. cisternarum

The transcriptome assembly process generated 1,350,426 contigs with median length of 329 bp (N50 = 668 bp) (Suppl. Table 1 and 2). Our BLAST survey of this assembly using *D. rerio* query sequences uncovered 11 *P. cisternarum* opsins: three visual opsins (*rh1.1*, *rh1.2*, and *lws*), and eight non-visual opsins (*exorh*, *va1*, *va2*, *parietopsin*, *tmt3a*, *opn4m3*, *peropsin*, and *rgr1*). Twenty-three opsins were identified in *I. punctatus*, and no additional *P.*

cisternarum genes were detected when *I. punctatus* opsins were employed as query sequences. The entire *P. cisternarum rh1.1* gene (a single-exon gene) was contained in one contig. The other 10 *P. cisternarum* opsins were incomplete; each was represented by one or more contigs that included between 44 bp and 747 bp of coding sequence that could be aligned to all other opsins. Despite this length variation, the *P. cisternarum* opsins were placed into well-supported clades with *I. punctatus* orthologs (Fig. 2, Suppl. File 1).

The observation that only *rh1.1* was full-length and that introns were present in all *P. cisternarum* opsins other than *rh1.1* and *rh1.2* suggests that most transcripts are not functional in the specimen studied. However, it is possible that intron retention (IR) is a form of translation regulation providing the potential to generate opsin proteins rapidly in response to light stimulation by splicing pre-existing transcripts. Another possibility is that opsin IR in *P. cisternarum* influences eye development by generating opsin mRNAs with premature termination codons (PTCs). Opsin loss can lead to retinal degeneration in humans (Silverman et al., 2020), thus nonsense-mediated decay (NMD) of intron-containing *P. cisternarum* opsin transcripts might play a role in eye degeneration.

P. cisternarum retains a rudimentary lens and shows reduced hair cells

Our histological analysis showed that the small eye of *P. cisternarum* is composed of an RPE and a rudimentary lens, we did not observe the cornea or sclera and the retina layers were absent in all specimens analyzed in this study. The lack of photoreceptor cells is interesting considering that we found opsin expression in the transcriptome. In addition to the histological analysis of RPE, we found RPE transcripts encoding for proteins such as retinoid isomer hydrolase (RPE65), GTP cyclohydrolase 1-like, L-dopachrome tautomerase, RPE-retinal G protein-coupled receptor-like, among others, in the transcriptome (**Fig. 3A, Suppl. Table 3**). Immunostaining revealed that among the lens-specific genes identified in our transcriptome, gamma-crystallin, a typical lens protein, was mainly expressed in border of the lens (**Fig. 3B**). In addition, our transcriptome analysis showed the expression of 36 genes that are critical for lens development and maintenance. This suggests that *P. cisternarum* retained a rudimentary lens, and that ocular reduction might be due to lower levels of

expression of these genes, however, to test this hypothesis, we would need to perform additional gene expression analysis (Suppl. Table 4) (Alunni *et al.*, 2007; Atukorala and Franz-Odenaal, 2018; Cavalleiro *et al.*, 2017; Hooven *et al.*, 2004; Yamamoto *et al.*, 2004).

Gross *et al.*, have shown previously that a set of genes important for lens maintenance and development is exclusively expressed in *A. mexicanus* surface population (lost in cave population). Comparative analysis of *A. mexicanus* natural populations identified 16 genes exclusively expressed in surface, but not cave populations (Gross *et al.*, 2013). Among those, we found 8 genes (*Rom1*, *Gnat1*, *Pde6*, *Crygm1*, *Crygm3*, *Lhx1b*, *Arr3a* and *Atp6ap1*) in the transcriptome of *P. cisternarum* (**Table 1, Suppl. Files 2-9**). These results suggest that the genetic mechanism of ocular reduction in *P. cisternarum* may be different from that described for *A. mexicanus*.

In cavefish, an increase in the number and size of neuromasts is one of the mechanisms used to offset eye degeneration (Soares and Niemiller, 2018; Van Trump *et al.*, 2010). Our results showed the presence of hair cells along the reduced lateral line and present in the head, however, not in abundance, as previously reported for *A. mexicanus* (Van Trump *et al.*, 2010) (**Fig. 3C,D**). The lack of supernumerary neuromasts in *P. cisternarum* could be linked to low water currents in these confined water streams of phreatic environments.

Conclusions

Previous studies in cavefish have uncovered genetic and developmental mechanisms underlying eye degeneration in dark-adapted fish. Here, we present evidence that eye degeneration in *P. cisternarum* might occur via mechanisms distinct from those described for cavefish species. Histological and molecular analyses in adult fish showed that *P. cisternarum* maintains eye structures such as RPE and a small lens. Our RNA-seq analyses revealed the visual and non-visual opsin repertoire in *P. cisternarum*, we identified a set of transcripts implicated in the development and maintenance of the RPE, retina as well as lens. Finally, adaptation to a subterranean environment in *P. cisternarum* appears to have occurred without concomitant enhancement of sensory neuromasts. In sum, our results provide the groundwork for future

studies aimed at identifying the genetic and developmental underpinnings of eye degeneration in *P. cisternarum* and dark-adapted species in general.

Ethics

This study was approved by IBAMA/SISBIO under license number 66015-1 and by the Ethics Committee for Animal Research at the Universidade Federal do Pará (protocol number 037-2015).

Data accessibility

Sequence data have been deposited in GenBank with the following BioProject accession number: PRJNA491408 and SRA file: SRR7878036.

Authors' contributions

PNS conceived the ideas and experimental design. LNP, BRM and JL collected the data; LNP, JST, RAPM and PNS analyzed the data; MPS and JT performed the transcriptome assembly, JT and AL performed opsin analysis; PNS, BRM and LNP led the writing of the manuscript. All authors contributed critically to the drafts, gave final approval for publication and agree to be held accountable for the work performed therein.

Acknowledgements

We especially thank Igor Schneider and Sylvain Darnet for insightful comments on this manuscript.

Conflict of Interest statement. None declared.

Funding

Funding from MCTIC/FINEP/FNDCT/AT Amazonia Legal to PNS. supported this work.

REFERENCES

- ALUNNI A, MENUET A, CANDAL E, PÉNIGAULT J-B, JEFFERY WR, RÉTAUX S (2007). Developmental mechanisms for retinal degeneration in the blind cavefish *Astyanax mexicanus*. *J Comp Neurol* 505: 221–233. Available at: <http://doi.wiley.com/10.1002/cne.21488>.
- ATUKORALA ADS, FRANZ-ODENDAAL TA (2018). Genetic linkage between altered tooth and eye development in lens-ablated *Astyanax mexicanus*. *Dev Biol* 441: 235–241. Available at: <https://doi.org/10.1016/j.snb.2019.127013>.
- BEAUDRY FEG, IWANICKI TW, MARILUZ BRZ, DARNET S, BRINKMANN H, SCHNEIDER P, TAYLOR JS (2017). The non-visual opsins: eighteen in the ancestor of vertebrates, astonishing increase in ray-finned fish, and loss in amniotes. *J Exp Zool Part B Mol Dev Evol* 328: 685–696.
- BOLGER AM, LOHSE M, USADEL B (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30: 2114–2120. Available at: <https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btu170>.
- CAVALHEIRO GR, MATOS-RODRIGUES GE, ZHAO Y, GOMES AL, ANAND D, PREDES D, DE LIMA S, ABREU JG, ZHENG D, LACHKE SA, CVEKL A, MARTINS RAP (2017). N-myc regulates growth and fiber cell differentiation in lens development. *Dev Biol* 429: 105–117. Available at: <https://doi.org/10.1016/j.snb.2019.127013>.
- DAVIES WIL, TAMAI TK, ZHENG L, FU JK, RIHEL J, FOSTER RG, WHITMORE D, HANKINS MW (2015). An extended family of novel vertebrate photopigments is widely expressed and displays a diversity of function. *Genome Res* 25: 1666–1679. Available at: <http://genome.cshlp.org/lookup/doi/10.1101/gr.189886.115>.
- FERNANDEZ L, SAUCEDO LJ, CARVAJAL-VALLEJOS FM, SCHAEFER SA (2007). A new phreatic catfish of the genus *Phreatobius* Goeldi 1905 from groundwaters of the Iténez River, Bolivia (Siluriformes: Heptapteridae). *Zootaxa*: 51–58.
- GASTEIGER E, HOOGLAND C, GATTIKER A, DUVAUD S, WILKINS MR, APPEL RD, BAIROCH A (2005). Protein Identification and Analysis Tools on the ExPASy Server. In *The Proteomics Protocols Handbook* Humana Press, Totowa, NJ, pp. 571–607. Available at:

- <http://link.springer.com/10.1385/1-59259-890-0:571>.
- GRABHERR MG, HAAS BJ, YASSOUR M, LEVIN JZ, THOMPSON DA, AMIT I, ADICONIS X, FAN L, RAYCHOWDHURY R, ZENG Q, et al. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol* 29: 644–652. Available at: <http://www.nature.com/articles/nbt.1883>.
- GROSS JB, FURTERER A, CARLSON BM, STAHL BA (2013). An Integrated Transcriptome-Wide Analysis of Cave and Surface Dwelling *Astyanax mexicanus* Ed. D Soares. *PLoS One* 8: e55659. Available at: <https://dx.plos.org/10.1371/journal.pone.0055659>.
- HALL TA (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41: 95–98. Available at: <http://jwbrown.mbio.ncsu.edu/JWB/papers/1999Hall1.pdf>.
- HOOVEN TA, YAMAMOTO Y, JEFFERY WR (2004). Blind cavefish and heat shock protein chaperones: a novel role for hsp90alpha in lens apoptosis. *Int J Dev Biol* 48: 731–738. Available at: <http://www.intjdevbiol.com/paper.php?doi=041874th>.
- JEFFERY WR (2009). Evolution and development in the cavefish *Astyanax*. *Curr Top Dev Biol* 86: 191–221.
- KIERNAN JA (2008). *Histological and histochemical methods: Theory and Practice*, 4th ed. Scion Publishing Limited.
- KUMAR S, STECHER G, TAMURA K (2016). MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33: 1870–1874. Available at: <https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msw054>.
- LANGMEAD B, SALZBERG SL (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9: 357–359. Available at: <http://www.nature.com/articles/nmeth.1923>.
- LI H, HANDSAKER B, WYSOKER A, FENNELL T, RUAN J, HOMER N, MARTH G, ABECASIS G, DURBIN R (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078–2079. Available at: <https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btp352>.

- LIN J-J, WANG F-Y, LI W-H, WANG T-Y (2017). The rises and falls of opsin genes in 59 ray-finned fish genomes and their implications for environmental adaptation. *Sci Rep* 7: 15568.
- MA L, NG M, WEELE CM, YOSHIZAWA M, JEFFERY WR (2020). Dual roles of the retinal pigment epithelium and lens in cavefish eye degeneration. *J Exp Zool Part B Mol Dev Evol*: jez.b.22923. Available at: <https://onlinelibrary.wiley.com/doi/abs/10.1002/jez.b.22923>.
- MENG F, ZHAO Y, POSTLETHWAIT JH, ZHANG C (2013). Differentially-expressed opsin genes identified in *Sinocyclocheilus* cavefish endemic to China. *Curr Zool* 59: 170–174. Available at: <https://academic.oup.com/cz/article-lookup/doi/10.1093/czoolo/59.2.170>.
- MILNE I, STEPHEN G, BAYER M, COCK PJA, PRITCHARD L, CARDLE L, SHAW PD, MARSHALL D (2013). Using Tablet for visual exploration of second-generation sequencing data. *Brief Bioinform* 14: 193–202. Available at: <https://academic.oup.com/bib/article-lookup/doi/10.1093/bib/bbs012>.
- MOHUN SM, DAVIES WL, BOWMAKER JK, PISANI D, HIMSTEDT W, GOWER DJ, HUNT DM, WILKINSON M (2010). Identification and characterization of visual pigments in caecilians (Amphibia: Gymnophiona), an order of limbless vertebrates with rudimentary eyes. *J Exp Biol* 213: 3586–3592. Available at: <http://jeb.biologists.org/cgi/doi/10.1242/jeb.045914>.
- MURIEL-CUNHA J, PINNA M de (2005). New data on cistern catfish, *Phreatobius cisternarum*, from subterranean waters at the mouth of the Amazon River (Siluriformes, Incertae Sedis). *Papéis Avulsos Zool (São Paulo)* 45: 327–339. Available at: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0031-10492005002600001&lng=en&nrm=iso&tlng=en.
- OHARA WM, DA COSTA ID, FONSECA ML (2016). Behaviour, feeding habits and ecology of the blind catfish *Phreatobius sanguijuela* (Ostariophysi: Siluriformes). *J Fish Biol* 89: 1285–1301. Available at: <http://doi.wiley.com/10.1111/jfb.13037>.
- OWENS GL, WINDSOR DJ, MUI J, TAYLOR JS (2009). A Fish Eye Out of Water: Ten Visual Opsins in the Four-Eyed Fish, *Anableps anableps* Ed.

- J Welch. *PLoS One* 4: e5970. Available at:
<https://dx.plos.org/10.1371/journal.pone.0005970>.
- PARTHA R, CHAUHAN BK, FERREIRA Z, ROBINSON JD, LATHROP K, NISCHAL KK, CHIKINA M, CLARK NL (2017). Subterranean mammals show convergent regression in ocular genes and enhancers, along with adaptation to tunneling. *Elife* 6: 1–26.
- PERTEA M, PERTEA GM, ANTONESCU CM, CHANG T-C, MENDELL JT, SALZBERG SL (2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol* 33: 290–295.
Available at: <http://www.nature.com/articles/nbt.3122>.
- PRIYAM A, WOODCROFT BJ, RAI V, MOGHUL I, MUNAGALA A, TER F, CHOWDHARY H, PIENIAK I, MAYNARD LJ, GIBBINS MA, et al. (2019). Sequenceserver: A Modern Graphical User Interface for Custom BLAST Databases. *Mol Biol Evol* 36: 2922–2924. Available at:
<https://academic.oup.com/mbe/article/36/12/2922/5549819>.
- PROTAS M, JEFFERY WR (2012). Evolution and development in cave animals: from fish to crustaceans. *Wiley Interdiscip Rev Dev Biol* 1: 823–845. Available at: <http://doi.wiley.com/10.1002/wdev.61>.
- RENNISON DJ, OWENS GL, TAYLOR JS (2012). Opsin gene duplication and divergence in ray-finned fish. *Mol Phylogenet Evol* 62: 986–1008.
- SCHOLTYSSEK C, KELBER A (2017). Farbensehen der Tiere Color vision in animals. *Der Ophthalmol* 114: 978–985.
- SEPPEY M, MANNI M, ZDOBNOV EM (2019). BUSCO: Assessing Genome Assembly and Annotation Completeness. In *Nagapattinam to Suvarnadwipa* pp. 227–245. Available at:
http://link.springer.com/10.1007/978-1-4939-9173-0_14.
- SHIBATTA OA, MURIEL-CUNHA J, PINNA MCC de (2007). A new subterranean species of Phreatobius Goeldi, 1905 (Siluriformes, Incertae sedis) from the Southwestern Amazon basin. *Pap Avulsos Zool* 47: 191–201. Available at:
http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0031-10492007001700001&lng=en&tlng=en.
- SOARES D, NIEMILLER ML (2018). Extreme Adaptation in Caves. *Anat Rec* 303: 15–23. Available at:

<https://onlinelibrary.wiley.com/doi/abs/10.1002/ar.24044>.

STEMMER M, SCHUHMACHER L-N, FOULKES NS, BERTOLUCCI C, WITTBRODT J (2015). Cavefish eye loss in response to an early block in retinal differentiation progression. *Development* 142: 743–752. Available at: <http://dev.biologists.org/cgi/doi/10.1242/dev.114629>.

THOMPSON JD, HIGGINS DG, GIBSON TJ (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22: 4673–4680. Available at: <https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/22.22.4673>.

TRAJANO E (2007). The Challenge of Estimating the Age of Subterranean Lineages: Examples from Brazil. *Acta Carsologica* 36: 191–198. Available at: <http://ojs.zrc-sazu.si/carsologica/article/view/221>.

VAN TRUMP WJ, COOMBS S, DUNCAN K, MCHENRY MJ (2010). Gentamicin is ototoxic to all hair cells in the fish lateral line system. *Hear Res* 261: 42–50. Available at: <https://linkinghub.elsevier.com/retrieve/pii/S0378595510000031>.

YAMAMOTO Y, STOCK DW, JEFFERY WR (2004). Hedgehog signalling controls eye degeneration in blind cavefish. *Nature* 431: 844–847. Available at: <http://www.nature.com/articles/nature02864>.

YOSHIZAWA M, GORIČKI Š, SOARES D, JEFFERY WR (2010). Evolution of a Behavioral Shift Mediated by Superficial Neuromasts Helps Cavefish Find Food in Darkness. *Curr Biol* 20: 1631–1636. Available at: <https://linkinghub.elsevier.com/retrieve/pii/S0960982210009164>.




Fig. 1. The *Phreatobius cisternarum* characterization. *P. cisternarum* adult specimen with miniaturization body, a slender head, and six long thin barbels **(A)**. Map showing collecting site of *P. cisternarum* in Benevides-Para-Brazil (the yellow circle indicates the Benevides city) **(B)**. Scale bar: 0.5 cm.




Fig. 2. Phylogenetic tree showing the visual and non-visual opsins of *Phreatobius cisternarum* (in red), *Danio rerio* and *Ictalurus punctatus*. Maximum parsimony tree showing relationships among *Phreatobius cisternarum*, *Ictalurus punctatus* and *Danio rerio* opsins. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown. Sequences were truncated at the 5' and 3' ends as these regions could not be aligned for all opsins with confidence. *rh1.2* and *va1* sequences from *P. cisternarum* clustered with orthologs from *I. punctatus*, but for these two genes the truncated alignment (Table S1) contained insufficient data to show their respective relationships to *D. rerio* orthologs. Monophyly of *rh1.2* and *va1* clades was well supported in analyses of longer subfamily-only alignments (not shown). *Danio rerio* has 42 opsins (Davies et al., 2015) and *I. punctatus* has 23 opsins. All were included in a preliminary analysis but only orthologs of *P. cisternarum* opsins were included in these analysis.




Fig. 3. The *Phreatobius cisternarum* remaining eye structures. Hematoxylin and eosin staining of head sections of *P. cisternarum* showing a retinal pigment epithelium (RPE) and lens (**A**). Immunofluorescence assay showing gamma crystallin expression on the lens (green) and DNA staining with DAPI (blue) (**B**). The distribution of superficial neuromasts (yellow dots) on the anterior lateral line (head) (**C**) and posterior lateral line (trunk) (**D**) of *P. cisternarum* specimens with DASPEI staining. Cryosections are 20 μm thick. Scale bars: 200 μm (**A** and **B**) and 2 mm (**C** and **D**).

Table 1. Lens genes expressed in the surface population of *A. mexicanus* and RNASeq from the head of *Phreatobius cisternarum*.

ID	Gene symbol	Gene name	TPM
TRINITY_DN133315_c2_g7_i1	Rom1	Retinal outer segment membrane protein 1	0,11575
TRINITY_DN145112_c3_g7_i3	Gnat1	Guanine nucleotide-binding protein (G protein), alpha transducing activity polypeptide 1	0,30194
TRINITY_DN138747_c6_g1_i2	Pde6	Phosphodiesterase 6G, cGMP-specific, rod	0,24273
TRINITY_DN139252_c5_g3_i1	Crygm1	Crystallin, gamma M1	0,13917
TRINITY_DN304038_c0_g1_i1	Crygm3	Crystallin, gamma M3	0,06959
TRINITY_DN131980_c0_g1_i2	Lhx1b	LIM class homeodomain protein	0,21923
TRINITY_DN146039_c1_g1_i1	Arr3a	Arrestin 3, retinal (X-arrestin), like	0,08267
TRINITY_DN143563_c2_g1_i9	Atp6ap1	ATPase, H ⁺ transporting, lysosomal accessory protein 1	1,63144