

# **On the role of pleiotrophin and its receptors in development and angiogenesis**

EVANGELIA PAPADIMITRIOU\*, ELENI MOURKOIANNI, DESPOINA NTENEKOU,  
MAGDALINI CHRISTOPOULOU, MARINA KOUTSIOUMPA and MARGARITA LAMPROU

## **ACCEPTED MANUSCRIPT**

The International Journal of Developmental Biology publishes an “Accepted” manuscript format as a free service to authors in order to expedite the dissemination of scientific findings to the research community as soon as possible after acceptance following peer review and corresponding modification (where appropriate).

An “Accepted” manuscript is published online prior to copyediting, formatting for publication and author proofing, but is nonetheless, fully citable through its Digital Object Identifier (DOI). Nevertheless, this “Accepted” version is NOT the final version of the manuscript.

When the final version of this paper is published within a definitive issue of the journal with copyediting, full pagination, etc. the new final version will be accessible through the same doi and this "Accepted" version of the paper will disappear.

**On the role of pleiotrophin and its receptors in development and angiogenesis**

Evangelia Papadimitriou<sup>\*</sup>, Eleni Mourkogianni, Despoina Ntenekou, Magdalini Christopoulou,  
Marina Koutsioumpa<sup>#</sup>, Margarita Lamprou

Laboratory of Molecular Pharmacology, Department of Pharmacy, University of Patras,  
GR26504 Patras, Greece

<sup>#</sup>*Current address:* Center for Systems Biomedicine, Vatche and Tamar Manoukian Division of Digestive Diseases, David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA, 90095, USA

**Running title:** PTN in development and angiogenesis

**Key words:** angiogenesis, development, growth factors, pleiotrophin, receptors

**\*Corresponding author:** Evangelia Papadimitriou, Ph.D., Laboratory of Molecular Pharmacology, Department of Pharmacy, University of Patras, GR 26504 Patras, Greece. Tel: 0030-2610-962336, e-mail: [epapad@upatras.gr](mailto:epapad@upatras.gr)

## **Abstract**

The secreted growth factor pleiotrophin (PTN) is expressed in all species and is evolutionary highly conserved, suggesting that it is significant for the regulation of important processes. The observation that it is highly expressed at early stages during development and at embryonic progenitor cells, highlights a potential important contribution to development. There is ample evidence for the role of PTN in the development of the nervous system and hematopoiesis, some but still not conclusive evidence for its role in the skeletomuscular system and limited evidence for its role in the development of other organs. Studies on its role in the cardiovascular system and angiogenesis suggest that PTN has a significant regulatory effect by acting on endothelial cells, while its role in smooth or cardiac muscle cells' functions has not been studied. This review highlights what is known up to date on the role of PTN in the development of various organs and in angiogenesis. Wherever known, evidence on the crosstalk among the receptors that mediate PTN's functions is also quoted, highlighting the complex regulatory pathways that affect development and angiogenesis.

## Introduction

The secreted growth factor pleiotrophin (PTN) was initially isolated from bovine uterus and fetal mouse brain by different groups in 1989. It was given several different names but the one that dominated and is currently being universally used is PTN (Fig. 1).

PTN is highly conserved among species and shows high amino acid homology with midkine (MK), the other member of the same family of heparin-binding proteins. The mature peptide consists of 136 amino acids, following cleavage of the signal peptide, and contains 24% cationic amino acids mainly at the amino- and carboxy-terminals, as well as 10 cysteines involved in the formation of 5 disulfide bonds. PTN comprises of two  $\beta$ -structures linked by a flexible region. Each  $\beta$ -structure contains three antiparallel  $\beta$ -folds, homologous to the thrombospondin type I repeat (TSR-1) motif. The ends of the molecule have numerous basal amino acids and lack configuration. The carboxy terminus of PTN has been shown responsible for its interaction with chondroitin sulfate chains. The human *ptn* gene of PTN has been identified on chromosome 7, in arm q33, is at least 42 kb and contains 7 exons. The open reading frame includes 4 exons, mostly exons 3 and 4, and the borders between introns/exons are well maintained. Different PTN isoforms have been suggested but have not been studied. Despite high homology in the coding region, the 3' and 5' untranslated regions differ between species. The promoter of the human *ptn* gene contains response elements for myogenic differentiation factor 1, trihelix transcription factor GT-1, activator protein 1, serum, homeobox A5 and Sox10. Possible binding sites for nuclear receptor  $\kappa$ B, cAMP binding protein, serum response factor and retinoic acid have been suggested but not proved (revised in Papadimitriou et al., 2009; 2016). Related to the latter, experimental evidence suggests that PTN expression is not enhanced by retinoic acid (Li *et al.*, 1992) and this notion is supported by our data showing that all-trans retinoic acid inhibits PTN expression (Theodorakopoulou *et al.*, 2006).

## PTN receptors

PTN has been shown to act through numerous receptors (Fig. 2). The first molecule identified as a functional PTN receptor in brain neurons was N-syndecan or syndecan-3, SDC3), which was found to interact with PTN in a solid phase binding assay. Using a similar assay, PTN was also found to interact with SDC1 in a heparan sulfate-dependent manner. It was later shown that the chondroitin sulphate chains on SDC1 and SDC4 are involved in their interaction with PTN, while the core protein of SDC1 may also have a role. A cooperative action of both TSR-1 domains of PTN is required for its interaction with the heparan sulphate chains of SDC3 and the regulation of synaptic plasticity. Despite these studies showing interaction of PTN with different syndecans, only interaction with SDC3 has been linked to PTN functions, as discussed later (revised in Papadimitriou *et al.*, 2009; 2016; Pantazaka and Papadimitriou, 2014).

Another early identified PTN receptor in the nervous system is receptor protein tyrosine phosphatase zeta 1 (PTPRZ1). PTN binds to high and low-affinity sites that involve both the chondroitin sulphate chains and the protein core of PTPRZ1. The carboxy-terminal PTN domain seems to be involved in the interaction with the PTPRZ1 chondroitin sulphate structures, which differentiate during development or in pathologies (revised in Pantazaka and Papadimitriou, 2014; Papadimitriou *et al.*, 2016). Binding of PTN to PTPRZ1 increases tyrosine phosphorylation of numerous downstream molecules, such as c-Src and Fyn kinases,  $\beta$ -catenin,  $\beta$ -adducin,  $\alpha_v\beta_3$  integrin, focal adhesion kinase, phosphoinositide 3-kinase and activation of protein kinase C alpha, beta or delta, ERK1/2 kinases (reviewed in Papadimitriou *et al.*, 2009; 2016) and cyclin dependent kinase 5 (Lampropoulou *et al.*, 2018). It was initially suggested that binding of PTN to PTPRZ1 inhibits the phosphatase activity of the receptor through dimerization/oligomerization (revised in Papadimitriou *et al.*, 2016), and as an answer to the doubts related to how could the highly

negatively charged chondroitin sulphate moieties on PTPRZ1 allow dimerization, it has been more recently suggested that the positively charged ligand PTN may neutralize electrostatic repulsion between chondroitin sulphate chains, thus inducing PTPRZ1 clustering, an effect also observed after removal of the PTPRZ1 chondroitin sulphate chains (Kuboyama *et al.*, 2016). However, this point needs further studies and clarification since there is also the possibility that the PTPRZ1 downstream signalling results from the regulation of tyrosine kinase receptors autophosphorylation by PTPRZ1. Such a mechanism has been described for the neuregulin receptor ErbB4 that interacts with PTPRZ1 through postsynaptic density-95 and results in decreased tyrosine phosphorylation of ErbB4 (Fujikawa *et al.*, 2007). Similarly, PTPRZ1 has been shown to dephosphorylate anaplastic lymphoma kinase (ALK) autophosphorylation sites, an effect that may be inhibited upon PTN binding to PTPRZ1 or following PTPRZ1 deletion or tyrosine phosphatase inhibition, leading to ALK activation (Xia *et al.*, 2019; Ntenekou *et al.*, 2020).

ALK has been suggested as a PTN receptor in ligand-receptor binding studies performed in cell-free assays and in intact cells, in which PTN was shown to bind to the ALK extracellular domain. Although direct PTN binding to ALK has been questioned by numerous studies, ALK seems to be activated by PTN and to be involved in PTN's signaling and actions. It has been suggested that PTN may activate ALK through PTPRZ1 (Papadimitriou *et al.*, 2016; Xia *et al.*, 2019) but the exact pathway needs to be further studied.

Another cell surface protein that was early identified and has been discussed as a low affinity PTN receptor is nucleolin (NCL). NCL is a ubiquitous nucleolar protein that regulates several aspects of the DNA and RNA metabolism, ribosome assembly and the development of various tissues and organs. NCL is present on the surface of activated endothelial and cancer cells following  $\alpha_v\beta_3$  integrin activation by PTN. Cell surface NCL interacts with  $\alpha_v\beta_3$  and PTPRZ1 and seems to affect

both PTN signaling and its translocation to the nucleus (Koutsoumpa and Papadimitriou, 2014; Papadimitriou *et al.*, 2016) through mechanisms that remain unclear.

Integrins are heterodimeric cell-membrane receptors that mediate cell-cell interactions and adhesion between the extracellular matrix and the cytoskeleton and play important roles during development (Maartens and Brown, 2015). Integrins that have been characterized as PTN receptors are  $\alpha_v\beta_3$  on endothelial cells (Mikelis *et al.*, 2009) and  $\alpha_M\beta_2$  on leukocytes (Shen *et al.*, 2017), mediating functions of PTN related to angiogenesis and inflammatory responses, respectively. PTN interacts with the extracellular domain of the  $\beta_3$  integrin subunit through its carboxy-terminal domain (Mikelis *et al.*, 2011), and with the  $\alpha_5$ - $\beta_5$  loop of the  $\alpha_M$ I-domain through its amino-terminal domain (Feng *et al.*, 2021). In endothelial cells,  $\alpha_v\beta_3$  interacts with PTPRZ1 even in the absence of PTN and this interaction is not affected by PTN (Mikelis *et al.*, 2009 and Fig. 3), although PTN induces PTPRZ1-dependent  $\beta_3$  Tyr774 phosphorylation (Mikelis *et al.*, 2009).

Neuropilin-1, a receptor for semaphorins that has significant role(s) in embryonic development of the nervous and the vascular system (Jubb *et al.*, 2012), has been shown to bind PTN. This interaction requires the TSR-1 domains of PTN and leads to ERK1/2 activation and enhancement of endothelial cell migration (Elahouel *et al.*, 2015), but has not been further studied.

More recently, we have shown that PTN selectively binds to vascular endothelial growth factor receptor 2 (VEGFR2) in endothelial and other types of cells and inhibits vascular endothelial growth factor A (VEGFA)-induced VEGFR2 phosphorylation at Tyr1175 (Lamprou *et al.*, 2020).

## **Expression and functions of PTN and its receptors during development**

The highest PTN expression is undoubtedly observed in the nervous system, the pituitary, the heart, the eye, the placenta, the seminal tissue and the testis, the bladder, and the stomach. PTPRZ1 is significantly expressed in the nervous system, the skin, and the eye. They are both highly expressed in induced pluripotent stem cells and embryonic stem cells (Papadimitriou *et al.*, 2016), suggesting important role(s) in early development. However, only in few cases a causal relationship between PTN and its receptors has been linked to specific functions.

### *Nervous system*

Since its discovery, PTN has been implicated in brain development and maturation. During development, it is highly expressed in the nervous system at sites of active mitogenesis, initially in the developing neuroepithelium and later in both glial cells and neurons in developing axonal tracts. In zebrafish embryos, overexpression of PTN induces extensive neurite outgrowth with complicated branching patterns. In the perinatal period, it is expressed by neurons, astrocytes and oligodendrocytes and has been characterized as a potent stimulator of neurite outgrowth in central nervous system neurons (revised in Winkler and Yao, 2004; González-Castillo *et al.*, 2015) and in spiral ganglion neurons (Bertram *et al.*, 2019). A recent study showed that neural stem cells secrete PTN into the niche and this contributes to the newborn neurons' maturation (Tang *et al.*, 2019). In adults, PTN expression is limited to specific neuronal subpopulations in the brain cortex, the hippocampus, the cerebellum, and the olfactory bulb (González-Castillo *et al.*, 2015).

A specific role of PTN has been described in relation to dopaminergic neurons. It is highly expressed in neural stem cells of mouse ventral mesencephalon and promotes the production of



dopaminergic neurons (González-Castillo *et al.*, 2015). It has been also shown to induce the differentiation of human umbilical cord mesenchymal stem cells into dopaminergic neuron-like cells (Yang *et al.*, 2013), and in combination with stromal cell-derived factor 1, insulin-like growth factor 2, and ephrin B1, to promote the differentiation of human embryonic stem cells to functional tyrosine hydroxylase-positive neurons (Vazin *et al.*, 2009). PTN is overexpressed in neurodegenerative diseases and confers a protective and/or nourishing effect on dopaminergic neurons *in vivo* and *in vitro* (Mourlevat *et al.*, 2005). The effects of PTN on the development of the nigrostriatal dopaminergic pathway are shown to be mediated by SDC3 and PTPRZ1 (González-Castillo *et al.*, 2015).

PTPRZ1 during brain development is expressed mainly to the ventricular and subventricular zone (Levy *et al.*, 1993), on both neuronal and glial cells (Canoll *et al.*, 1993). *Ptprz1*-knockout mice show neurological abnormalities and differences in behavior and learning, including increased exploratory activities to novelty, deficits in spatial and contextual learning, and reduced responses to methamphetamine (Tanga *et al.*, 2019). PTN also maintains the self-renewing phenotype of oligodendrocyte progenitor cells through PTPRZ1 (McClain *et al.*, 2012) that has been shown to be important for the survival and recovery of oligodendrocytes from demyelinating disease (Harroch *et al.*, 2002). More recently, PTPRZ1 has been shown to be important for the perineuronal net structures that are important for both development and plasticity of the brain in the adult (Eill *et al.*, 2020).

PTN has been shown to mediate the neurite stimulating activity of the peptide Y-P30 through binding to SCD2 and SDC3 (Landgraf *et al.*, 2008). Phosphacan, the alternatively spliced extracellular PTPRZ1 domain, inhibits neurite outgrowth (Margolis *et al.*, 1996), and PTN has been shown to convert the neurite growth inhibitory effect of chondroitin sulfate proteoglycans

into an activating effect (Paveliev *et al.*, 2016). The expression pattern of PTN, PTPRZ1 and SDC3 in the postnatal mouse cerebellum supports their implication in the morphogenesis of Purkinje cells, as well as in the control of the granule cell migration (Basille-Dugay *et al.*, 2013), while disruption of PTN distribution extracellularly alters the development and function of the neuronal circuits of the cerebellum (Hamza *et al.*, 2016). Besides central nervous system, PTN is highly expressed in the peripheral nervous system during development and seems to promote the repair of neurons after injury of peripheral nerves. It may also have a significant role in muscle innervation, based on the observations that it is present at the neuromuscular junction and contributes to acetylcholine receptors clustering (revised in Jin *et al.*, 2009).

The effects of PTN on neural development and neurite outgrowth have been shown to be also mediated by ALK (Yanagisawa *et al.*, 2010). ALK has been implicated in the nervous system development, signaling during neuromuscular junction development in *C. elegans*, affecting the establishment of iridophores and normal pigmentation patterns in zebrafish, and neurogenesis in mice. It is highly expressed during embryonic mouse and chick development, in both the central and peripheral nervous system, while ALK expression levels are decreased after birth (revised in Kalamatianos *et al.*, 2018). Whether any of these functions are related to PTN has not been studied.

### *Muscle*

Outside the nervous system, early after its discovery, PTN was found to be expressed in human intestinal smooth muscle cells (Li *et al.*, 1992) and in satellite cells (Wanaka *et al.*, 1993). However, although a potential role of PTN in myogenesis was early suspected, the first evidence came from studies showing that PTN mRNA and protein expression are increased during

myogenesis and regeneration after crushing, in newly formed myotubes and in activated myoblasts before fusion *in vivo*. *In vitro*, PTN expression increases during the differentiation process, being maximal on fusion of myoblasts into myotubes (Caruelle *et al.*, 2004).

PTN expression is found increased during postnatal heart development and PTN increases postnatal cardiomyocytes' DNA synthesis (Chen *et al.*, 2004), suggesting that PTN may have a role in myocardial development and regeneration. PTN overexpression in slow-twitch soleus muscle leads to an increased number of pure type 1 fibers and decreased number of pure type 2A and 2X fibers, as well as increased sulfonylurea receptor 1, citrate synthase and cyclooxygenase IV gene expression and increased vascularization, suggesting a shift toward a more oxidative metabolism and improved muscle performance (Camerino *et al.*, 2013). The role of PTN receptors in heart development has not been studied. We have unpublished evidence using two different animal models, as well as human data showing that PTPRZ1 is expressed in the embryonic but not in the adult heart and plays a role in heart morphogenesis and subsequent function (Katraki-Pavlou *et al.*, unpublished).

As also mentioned above, PTN is a muscle protein present at the neuromuscular junction in close contact with the acetylcholine receptors and may have a significant role in muscle innervation (Jin *et al.*, 2009).

### *Skeletal system*

A role of PTN in bone development has been early suggested based on studies showing expression of its mRNA during bone growth in ancestral cartilage and pulp. PTN has been initially considered important for new bone formation based on observations that it induces osteoblast attachment to

the extracellular bone material through binding to a specific receptor, which maybe SDC3. It was also shown to induce hypertrophy during the chondrogenetic differentiation of human bone marrow progenitor cells, an effect that has been associated with both ALK and PTPRZ1. PTN enhances the release of heparin-binding epidermal growth factor, which then activates its receptor in precursor osteoblasts and osteoblast-type cells, increases alkaline phosphatase activity and inhibits dexamethasone-induced cell death. PTN overexpressing transgenic mice develop a phenotype characterized by increased bone growth and higher salt and bone density. When PTN is overexpressed solely in the mouse skeletal system, increased bone length and maturation are observed at the early stages of bone development, a phenomenon balanced in adult life. PTN-deficient mice appear to have bone growth retardation at two months, and osteopenia and resistance to bone remodeling in adult life (revised in Lamprou *et al.*, 2014). In transgenic mice that overexpress PTN, bone loss due to estrogen deficiency (Masuda *et al.*, 1997) or almost zero gravity (Tavella *et al.*, 2012) is compensated, in line with the observation that estrogen enhances PTN expression (Xi *et al.*, 2020). The increased expression of PTN and PTPRZ1 in osteoblasts depleted of insulin-like growth factor binding protein-2 has been suggested to help maintain, at least partly, normal bone mass in female mice (Xi *et al.*, 2020).

PTN is also involved in odontogenesis. It was initially shown that PTN is expressed during initiation, morphogenesis and cytodifferentiation stages of incisor development at structures that are important for tooth morphogenesis (Mitsiadis *et al.*, 2008). PTN is also expressed in epithelial and mesenchymal dental cell lines and its expression is regulated by bone morphogenetic proteins. During maturation of the ameloblasts and odontoblasts, PTN is expressed in the inner enamel epithelium; PTN is also expressed in the terminally differentiated and enamel matrix-secreting ameloblasts and odontoblasts of the adult mouse incisors and molars (Erlandsen *et al.*, 2012). More

recently, it has been shown that PTN positively regulates dental pulp stem cells proliferation and potential to differentiate (Jin *et al.*, 2020), protecting them from senescence (Zhang *et al.*, 2021).

### *Hematopoiesis and immune system*

PTN promotes hematopoietic stem cells (HSCs) expansion *in vitro* and HSCs regeneration *in vivo* (Himburg *et al.*, 2010) via PTPRZ1-mediated activation of the RAS pathway (Himburg *et al.*, 2014), supporting a role of the PTN/PTPRZ1 axis in hemopoiesis. The main source of PTN during steady-state hematopoiesis is the leptin receptor expressing bone marrow stromal cells, but regeneration of HSCs depends on endothelial cell-derived PTN (Himburg *et al.*, 2018). In the same line, PTN secreted into the human tonsil mesenchymal stem cells conditioned medium increases HSC transplantation efficacy (Kim *et al.*, 2020). It has been also shown that loss of stromal PTN results in changes in expression of genes that lead to myeloid engraftment dominance, suggesting a role of PTN in maintaining the balance of myeloid and lymphoid potential of regenerating HSCs (Istvanffy *et al.*, 2011).

PTN is mitogenic for human peripheral blood mononuclear cells and this initial observation suggested that it may be involved in the growth regulation of cell mediated immunity (Achour *et al.*, 2001), further supported by a subsequent study showing enhanced expression of inflammatory cytokines, such as tumor necrosis factor  $\alpha$  and interleukins 1 $\beta$  and 6 by PTN (Achour *et al.*, 2008). A role of PTN in adaptive immunity was corroborated by the observation that interferon  $\gamma$  induces PTN expression by macrophages (Li *et al.*, 2010) and a role in innate immunity has been suggested by its bactericidal properties (Svensson *et al.*, 2010; Guyot *et al.*, 2016). PTN has been also suggested as a potent regulator of neuroinflammation through PTPRZ1 (Herradon *et al.*, 2019),

and a regulator of the levels of functional cytokines in both M1 and M2 type microglia cells, strengthening M1/M2 transformation (Miao *et al.*, 2019).

Mice that are knockout for PTPRZ1 have a higher number and proportion of mature B cells and signaling of MK through PTPRZ1 leads to B cell survival (Cohen *et al.*, 2012). Whether PTN has such an effect on B cells has not been studied. It has been only shown that PTN expression is significantly increased in B cells from both chronic and acute leukemia patients compared to healthy controls and suppression of PTN activity induced apoptosis of B cells from both leukemia patients and cell lines (Du *et al.*, 2014).

### *Urogenital system*

In the human mammary gland, PTN is expressed in epithelial and endothelial cells (Ledoux *et al.*, 1997) and helps maintain the mammary epithelial cells in a progenitor phenotype, thus delaying the mammary gland maturation (Rosenfield *et al.*, 2012). Its expression in the uterus depends on the estrous cycle and is upregulated by progesterone (Milhiet *et al.*, 1998). In the endometrial stromal cells, it is up-regulated during decidualization (Bany and Schultz, 2001), apparently having a vital role in the progesterone-induced decidualization pathway (Yu *et al.*, 2018). High mobility group box 3 regulates uterine stromal cells proliferation and differentiation by targeting PTN (Wang *et al.*, 2019), further supporting a physiological role of PTN in uterine decidualization. CD49a<sup>+</sup> Eomes<sup>+</sup> natural killer (NK) cells in the human and mouse uterus secrete PTN and enhance fetal growth during early stages of fetal development (Fu *et al.*, 2017). The CD49a<sup>+</sup> PBX homeobox 1 enhances PTN transcription in decidual NK cells, thus promoting fetal

growth (Zhou *et al.*, 2020). Female mice that are deficient in both PTN and MK show reproductive abnormalities (Muramatsu *et al.*, 2006).

PTN is expressed in testicular Leydig cells and plays a normal role in spermatogenesis. Male mice deficient in PTN are characterized by infertility, atrophic testes and apoptotic sperm cells (Zhang *et al.*, 1999). PTN mRNA and protein are expressed in ventral mesenchymal pad and prostatic mesenchyme surrounding ductal epithelial tips that undergo branching morphogenesis *in vivo*. *In vitro*, PTN is up regulated by androgens and stimulates both stromal and epithelial cell proliferation, as well as branching morphogenesis (Orr *et al.*, 2011).

In the embryonic kidney, PTN is expressed in the developing kidney mesenchyme (Mitsiadis *et al.*, 1995) and localizes onto the basement membrane of the developing ureteric bud (Sakurai *et al.*, 2001).

### *Respiratory system*

PTN and PTPRZ1 have been shown to be highly expressed in mesenchymal and epithelial cells of fetal lungs, respectively. PTN promotes fetal type II cell proliferation and inhibits their trans-differentiation into alveolar epithelial type I cells, through multiple signaling pathways that include  $\beta$ -catenin and Notch (Weng *et al.*, 2009).

### *Eye*

PTN is mitogenic for bovine lens epithelial cells (Delbe *et al.*, 1995), and blocks or promotes rod or bipolar cell differentiation respectively (Roger *et al.*, 2006). In *Drosophila*, overexpression of

the MK/PTN orthologue *miple* leads to defective eye patterning through *Ptp99A*, the *Drosophila* ortholog of *PTPRZ1* (Muñoz-Soriano *et al.*, 2013). During embryonic development, PTN is expressed in the iris mesenchyme, the optic nerve, and the neural retina, suggesting that it may have a role in epithelial-mesenchymal interactions and affect optic nerve and retinal development. Among PTN receptors, *SDC1*, and *SDC3* are expressed in the cornea, whereas all receptors including *RPTPZ1* and *ALK* are expressed in the retina (Cui and Lwigale, 2019) but a causal relation to the PTN effects on the eye has not been studied.

### *Liver*

PTN was initially identified as a mitogen for adult hepatocytes (Sato *et al.*, 1999) and parenchymal cells in both adult and embryonic liver (Asahina *et al.*, 2002). It is expressed by mesenchymal cells in fetal liver, its expression is gradually reduced during development (Asahina *et al.*, 2002; Ito *et al.*, 2014) and seems to play a role in liver regeneration (Asahina *et al.*, 2002; Michelotti *et al.*, 2016) through its receptor *PTPRZ1* (Michelotti *et al.*, 2016). It was recently shown that in myofibroblasts there is decreased expression of several growth factors, among which PTN, compared to fibroblasts, and this leads to reduced hepatoblasts proliferation and induced maturation of the differentiated cholangiocytes (Wang *et al.*, 2020).

### *Other organs*

PTN expression has been identified during all embryonic stages in the neurohypophysis primordium. It is expressed only in neurohypophyseal cells and might be involved in pituitary development (Fujiwara *et al.*, 2014).



PTN mRNA is strongly expressed in the mouse cochlea one week after birth and gradually decreases, being undetectable by week 8 after birth. PTN knockout mice exhibit low to moderate auditory deficits and have higher hearing thresholds compared to the wild-type mice, with an almost normal appearance of the stria vascularis but weak expression of the Kir4.1 potassium channel (Zou et al., 2006). Mice knockout for both PTN and MK also appear with severe vacuolar degeneration in the intermediate cells (Sone *et al.*, 2011).

### **PTN and angiogenesis**

The vascular system is very important for fetal development and there is evidence that PTN has a significant regulatory role in vascular homeostasis, although the exact pathways involved are still being explored.

The initial evidence came from *in vitro* studies showing that PTN is mitogenic for endothelial cells. The mitogenic effect of PTN was questioned in subsequent studies and it seems that it may be cell-type and cell-context specific or depend on the origin of the recombinant protein used (revised in Papadimitriou et al., 2004; 2009). PTN induces endothelial cell migration and differentiation into tubes in various substrates *in vitro*, through its receptor PTPRZ1 that forms a functional complex with  $\alpha_v\beta_3$  integrin on the surface of endothelial cells and leads to cell surface translocation of NCL (revised in Papadimitriou et al., 2009; 2016). PTN has been also shown to enhance VEGFA expression (Koutsoumpa *et al.*, 2012; Palmieri *et al.*, 2015) and this may contribute to its angiogenic activities, although the receptors/pathways involved in this action have not been studied. In contrast to the initial characterization of PTN as an angiogenesis-promoting

growth factor, later studies suggested that PTN may also limit the angiogenic effect of VEGFA (Heroult *et al.*, 2004; Koutsoumpa *et al.*, 2015). The latter was further supported by data showing that VEGFA competes with PTN for binding to PTPRZ1 (Koutsoumpa *et al.*, 2015). Binding of VEGFA to PTPRZ1 inhibits endogenous PTN levels (Poimenidi *et al.*, 2016) that limit the angiogenic effects of VEGFA, thus enhancing VEGFA-induced endothelial cell migration (Koutsoumpa *et al.*, 2015; Poimenidi *et al.*, 2016). More recently, we have shown that PTN selectively binds VEGFR2 and inhibits its activation by VEGFA, highlighting another mechanism through which PTN limits the angiogenic effect of VEGFA (Lamprou *et al.*, 2020). Although the cross talk among PTPRZ1,  $\alpha_v\beta_3$  and VEGFR2 is still being studied, our up-to-date data support a model through which PTN has a moderate stimulatory effect on endothelial cells via PTPRZ1/ $\alpha_v\beta_3$ /cell surface NCL but limits the angiogenic effects of VEGFA through VEGFR2 (Fig. 4).

In line with a balancing effect on angiogenesis, PTN has moderate effects in *in vivo* animal models of angiogenesis, such as the rabbit cornea, the chorioallantoic membrane of the chick embryo, the ischemic myocardium in rats, a model of acute posterior muscle ischemia, slow-twitch soleus and fast-twitch extensor digitorum longus muscles, and numerous tumor models (revised in Papadimitriou *et al.*, 2009; 2016). When expressed by monocytes, it has been shown to lead to a decrease in the expression of monocyte markers and an increase in endothelial cell markers, inducing differentiation of monocytes into functional endothelial cells (Sharifi *et al.*, 2006). A role of PTN in angiogenesis is also supported by the expression of PTN and its receptors PTPRZ1, ALK, SDC1 and SDC3 in mesenchymal cells, fetal macrophages, and fetal vessels, in line with increasing levels of PTN in maternal bloodstream as pregnancy progresses (Ball *et al.*, 2009). PTN has been also suggested as a promising factor for inducing angiogenesis during aging based on the

observation that it restores the age-related attenuation of angiogenesis in the aortic ring model (Besse *et al.*, 2013).

The role of PTPRZ1 on angiogenesis has been mostly discussed based on its function as a PTN and VEGFA receptor, as mentioned above. We have recently shown that PTPRZ1 knockout mice show enhanced angiogenesis in several organs, such as lung, heart and the retina and microvascular endothelial cells isolated from the lungs of these animals have enhanced angiogenic properties (Ntenekou *et al.*, 2020; Katraki-Pavlou *et al.*, unpublished).

### **Conclusions**

PTN is a highly conserved, ubiquitously expressed protein that seems to play a notable role in the nervous system development and maturation. Data from several other organs show that it is also involved in epithelial-mesenchymal interactions during development and is expressed by endothelial cells, regulating angiogenesis. Due to the high homology degree with MK, the one compensates for the other in several but not all cases, and this adds to the complexity of studying PTN. Another point of complexity is the increasing number of receptors that are being identified to mediate the effects of PTN in several different types of cells and organs, forcing the researchers to try to clarify the interactions and the crosstalk of the different receptors/pathways involved in each case. Moreover, the PTN receptors identified up to date are not specific for PTN and mediate the effects of several other molecules besides PTN and MK (revised in Pantazaka and Papadimitriou, 2014; Papadimitriou *et al.*, 2016), suggesting that PTN may regulate the effects of other growth factors, as has been shown for VEGFA in endothelial cells (Koutsoumpa *et al.*, 2015; Poimenidi *et al.*, 2016). Although PTN does not seem to be indispensable for survival, it seems to

have important roles in several pathological conditions, affecting pathways that are deranged in pathological settings and this is still an open field to study. Meticulous *in vitro* work combined with high throughput assays and loss- or gain-of-function *in vivo* studies, will shed light on many unidentified aspects related to the role of PTN in development, angiogenesis, and pathologies.

### **Acknowledgements**

Eleni Mourkogianni and Despoina Ntenekou are recipients of PhD scholarships financed by the State Scholarship Foundation in Greece (IKY) (Operational Program “Human Resources Development – Education and Lifelong Learning”, Partnership Agreement 2014-2020).

## References

- ACHOUR A, LAAROUBI D, CARUELLE D, BARRITAU D, COURTY J (2001). The angiogenic factor heparin affinity regulatory peptide (HARP) induces proliferation of human peripheral blood mononuclear cells. *Cell Mol Biol (Noisy-le-grand)* 47 Online Pub: OL73-7.
- ACHOUR A, M'BIKA JP, BAUDOUIN F, CARUELLE D, COURTY J (2008). Pleiotrophin induces expression of inflammatory cytokines in peripheral blood mononuclear cells. *Biochimie* 90: 1791-1795.
- ASAHINA K, SATO H, YAMASAKI C, KATAOKA M, SHIOKAWA M, KATAYAMA S, TATENO C, YOSHIKATO K (2002). Pleiotrophin/heparin-binding growth-associated molecule as a mitogen of rat hepatocytes and its role in regeneration and development of liver. *Am J Pathol* 160: 2191-2205.
- BALL M, CARMODY M, WYNNE F, DOCKERY P, AIGNER A, CAMERON I, HIGGINS J, SMITH SD, APLIN JD, MOORE T. (2009). Expression of pleiotrophin and its receptors in human placenta suggests roles in trophoblast life cycle and angiogenesis. *Placenta* 30: 649-653.
- BANY BM, SCHULTZ GA (2001). Pleiotrophin messenger ribonucleic acid levels increase in mouse endometrial stromal cells during decidualization. *Endocrinology* 142: 955-958.
- BASILLE-DUGAY M, HAMZA MM, TASSERY C, PARENT B, RAOULT E, BÉNARD M, RAISMAN-VOZARI R, VAUDRY D, BUREL DC (2013). Spatio-temporal characterization of the pleiotrophinergic system in mouse cerebellum: Evidence for its key role during ontogenesis. *Exp Neurol* 247: 537-551.

BERTRAM S, ROLL L, REINHARD J, GROß K, DAZERT S, FAISSNER A, VOLKENSTEIN S (2019). Pleiotrophin increases neurite length and number of spiral ganglion neurons *in vitro*. *Exp Brain Res* 237: 2983-2993.

BESSE S, COMTE R, FRÉCHAULT S, COURTY J, JOËL DEL, DELBÉ J (2013). Pleiotrophin promotes capillary-like sprouting from senescent aortic rings. *Cytokine* 62: 44–47.

CAMERINO GM, PIERNO S, LIANTONIO A, DE BELLIS M, CANNONE M, SBLENDORIO V, CONTE E, MELE A, TRICARICO D, TAVELLA S, RUGGIU A, CANCEDDA R, OHIRA Y, DANIELI-BETTO D, CICILIOT S, GERMINARIO E, SANDONÀ D, BETTO R, CAMERINO DC, DESAPHY JF (2013). Effects of pleiotrophin overexpression on mouse skeletal muscles in normal loading and in actual and simulated microgravity. *PLoS One* 8: e72028.

CANOLL PD, BARNEA G, LEVY JB, SAP J, EHRLICH M, SILVENNOINEN O, SCHLESSINGER J, MUSACCHIO JM (1993). The expression of a novel receptor-type tyrosine phosphatase suggests a role in morphogenesis and plasticity of the nervous system. *Brain Res Dev Brain Res* 75: 293-298.

CARUELLE D, MAZOUZI Z, HUSMANN I, DELBÉ J, DUCHESNAY A, GAUTRON J, MARTELLY I, COURTY J (2004). Upregulation of HARP during *in vitro* myogenesis and rat soleus muscle regeneration. *J Muscle Res Cell Motil* 25: 45–53.

CHEN HW, YU SL, CHEN WJ, YANG PC, CHIEN CT, CHOU HY, LI HN, PECK K, HUANG CH, LIN F Y, CHEN JJ W, LEE YT (2004). Dynamic changes of gene expression profiles during postnatal development of the heart in mice. *Heart* 90: 927–934.

COHEN S, SHOSHANA OY, ZELMAN-TOISTER E, MAHARSHAK N, BINSKY-EHRENREICH I, GORDIN M, HAZAN-HALEVY I, HERISHANU Y, SHVIDEL L, HARAN

M, LENG L, BUCALA R, HARROCH S, SHACHAR I (2012). The cytokine midkine and its receptor RPTP $\zeta$  regulate B cell survival in a pathway induced by CD74. *J Immunol* 188: 259-269.

CUI R, LWIGALE P (2019). Expression of the heparin-binding growth factors Midkine and pleiotrophin during ocular development. *Gene Expr Patterns* 32: 28-37.

DELBÉ J, VACHEROT F, LAAROUBI K, BARRITAU D, COURTY J (1995). Effect of heparin on bovine epithelial lens cell proliferation induced by heparin affinity regulatory peptide. *J Cell Physiol* 164: 47-54.

DU CX, WANG L, LI Y, XIAO W, GUO QL, CHEN F, TAN XT (2014). Elevated expression of pleiotrophin in lymphocytic leukemia CD19+ B cells. *APMIS* 122: 905-13.

EILL GJ, SINHA A, MORAWSKI M, VIAPIANO MS, MATTHEWS RT (2020). The protein tyrosine phosphatase RPTP $\zeta$ /phosphacan is critical for perineuronal net structure. *J Biol Chem* 295: 955-68.

ELAHOUËL R, BLANC C, CARPENTIER G, FRECHAULT S, CASCONI I, DESTOUCHES D, DELBÉ J, COURTY J, HAMMA-KOURBALI Y (2015). Pleiotrophin exerts its migration and invasion effect through the neuropilin-1 pathway. *Neoplasia* 17: 613-624.

ERLANDSEN H, AME, JE, TAMKENATH A, MAMAEVA O, STIDHAM K, WILSON ME, PEREZ-PINERA P, DEUEL TF, MACDOUGALL M (2012). Pleiotrophin Expression during Odontogenesis. *J Histochem Cytochem* 60: 366–375.

FENG W, NGUYEN H, SHEN D, DENG H, JIANG Z, PODOLNIKOVA N, UGAROVA T, WANG X (2021). Structural Characterization of the Interaction between the  $\alpha$ MI-Domain of the Integrin Mac-1 ( $\alpha_M\beta_2$ ) and the Cytokine Pleiotrophin. *Biochemistry* 60: 182-193.

FU B, ZHOU Y, NI X, TONG X, XU X, DONG Z, SUN R, TIAN Z, WEI H (2017). Natural Killer Cells Promote Fetal Development through the Secretion of Growth-Promoting Factors. *Immunity* 47: 1100-1113.

FUJIKAWA A, CHOW JP, SHIMIZU H, FUKADA M, SUZUKI R, NODA M (2007). Tyrosine phosphorylation of ErbB4 is enhanced by PSD95 and repressed by protein tyrosine phosphatase receptor type Z. *J Biochem* 142: 343-350.

FUJIWARA K, MALIZA R, TOFRIZAL A, BATCHULUUN K, RAMADHANI D, TSUKADA T, AZUMA M, HORIGUCHI K, KIKUCHI M, YASHIRO T (2014). In situ hybridization analysis of the temporospatial expression of the midkine/pleiotrophin family in rat embryonic pituitary gland. *Cell Tissue Res* 357: 337–344.

GONZÁLEZ-CASTILLO C, ORTUÑO-SAHAGÚN D, GUZMÁN-BRAMBILA C, PALLÀS M, ROJAS-MAYORQUÍN AE (2015). Pleiotrophin as a central nervous system neuromodulator, evidences from the hippocampus. *Front Cell Neurosci* 8: 443.

GUYOT N, LABAS V, HARICHAUX G, CHESSÉ M, POIRIER JC, NYS Y, RÉHAULT-GODBERT S (2016). Proteomic analysis of egg white heparin-binding proteins: towards the identification of natural antibacterial molecules. *Sci Rep* 6: 27974.

HAMZA MM, REY SA, HILBER P, ARABO A, COLLIN T, VAUDRY D, BUREL D (2016). Early Disruption of Extracellular Pleiotrophin Distribution Alters Cerebellar Neuronal Circuit Development and Function. *Molecular Neurobiology* 53: 5203–5216.

HARROCH S, FURTADO GC, BRUECK W, ROSENBLUTH J, LAFAILLE J, CHAO M, BUXBAUM JD, SCHLESSINGER J (2002). A critical role for the protein tyrosine phosphatase receptor type Z in functional recovery from demyelinating lesions. *Nat Genet* 32: 411-4.



HÉROULT M, BERNARD-PIERROT I, DELBÉ J, HAMMA-KOURBALI Y, KATSORIS P, BARRITAU D, PAPADIMITRIOU E, PLOUET J, COURTY J (2004). Heparin affinity regulatory peptide binds to vascular endothelial growth factor (VEGF) and inhibits VEGF-induced angiogenesis. *Oncogene* 23: 1745–1753.

HERRADON G, RAMOS-ALVAREZ MP, GRAMAGE E (2019). Connecting Metainflammation and Neuroinflammation Through the PTN-MK-RPTP $\beta/\zeta$  Axis: Relevance in Therapeutic Development. *Front Pharmacol* 10: 377.

HIMBURG HA, MURAMOTO GG, DAHER P, MEADOWS SK, RUSSELL JL, DOAN P, CHI JT, SALTER AB, LENTO WE, REYA T, CHAO NJ, CHUTE JP (2010). Pleiotrophin regulates the expansion and regeneration of hematopoietic stem cells. *Nat Med* 16: 475-482.

HIMBURG HA, TERMINI CM, SCHLUSSEL L, KAN J, LI M, ZHAO L, FANG T, SASINE JP, CHANG VY, CHUTE JP (2018). Distinct Bone Marrow Sources of Pleiotrophin Control Hematopoietic Stem Cell Maintenance and Regeneration. *Cell Stem Cell* 23: 370-381.e5.

HIMBURG HA, YAN X, DOAN PL, QUARMYNE M, MICEWICZ E, MCBRIDE W, CHAO NJ, SLAMON DJ, CHUTE JP (2014). Pleiotrophin mediates hematopoietic regeneration via activation of RAS. *J Clin Invest* 124: 4753-4758.

ISTVANFFY R, KRÖGER M, ECKL C, GITZELMANN S, VILNE B, BOCK F, GRAF S, SCHIEMANN M, KELLER UB, PESCHEL C, OOSTENDORP RA (2011). Stromal pleiotrophin regulates repopulation behavior of hematopoietic stem cells. *Blood* 118: 2712-2722.

ITO K, YANAGIDA A, OKADA K, YAMAZAKI Y, NAKAUCHI H, KAMIYA A (2014). Mesenchymal progenitor cells in mouse foetal liver regulate differentiation and proliferation of hepatoblasts. *Liver Int* 34: 1378–1390.

JIN L, GAO F, ZHANG L, WANG C, HU L, FAN Z, XIA D (2020). Pleiotropin enhances the osteo/dentinogenic differentiation potential of dental pulp stem cells. *Connect Tissue Res* 24: 1-13.

JIN L, JIANGHAI C, JUAN L, HAO K (2009). Pleiotrophin and peripheral nerve injury. *Neurosurg Rev* 32: 387-393.

JUBB AM, STRICKLAND LA, LIU SD, MAK J, SCHMIDT M, KOEPPEN H (2012). Neuropilin-1 expression in cancer and development. *J Pathol* 226: 50-60.

KALAMATIANOS T, DENEKOU D, STRANJALIS G, PAPADIMITRIOU E (2018). Anaplastic Lymphoma Kinase in Glioblastoma: Detection/Diagnostic Methods and Therapeutic Options. *Recent Pat Anticancer Drug Discov* 13: 209-223.

KIM YH, CHO KA, LEE HJ, PARK M, SHIN SJ, PARK JW, WOO SY, RYU KH (2020). Conditioned Medium from Human Tonsil-Derived Mesenchymal Stem Cells Enhances Bone Marrow Engraftment via Endothelial Cell Restoration by Pleiotrophin. *Cells* 9: 221.

KOUTSIOUMPA M, DROSOU G, MIKELIS C, THEOCHARI K, VOURTSIS D, KATSORIS P, GIANNOPOULOU E, COURTY J, PETROU C, MAGAFA V, CORDOPATIS P, PAPADIMITRIOU E (2012). Pleiotrophin expression and role in physiological angiogenesis *in vivo*: Potential involvement of nucleolin. *Vasc Cell* 4: 1–13.

KOUTSIOUMPA M, PAPADIMITRIOU E (2014). Cell surface nucleolin as a target for anti-cancer therapies. *Recent Pat Anticancer Drug Discov* 9: 137-152.

KOUTSIOUMPA M, POIMENIDI E, PANTAZAKA E, THEODOROPOULOU C, SKOURA A, MEGALOOIKONOMOU V, KIEFFER N, COURTY J, MIZUMOTO S, SUGAHARA K,

PAPADIMITRIOU E (2015). Receptor protein tyrosine phosphatase beta/zeta is a functional binding partner for vascular endothelial growth factor. *Mol Cancer* 14: 1–16.

KOUTSIOUMPA M, POLYTARCHOU C, COURTY J, ZHANG Y, KIEFFER N, MIKELIS C, SKANDALIS SS, HELLMAN U, ILIOPOULOS D, PAPADIMITRIOU E (2013). Interplay between  $\alpha_v\beta_3$  integrin and nucleolin regulates human endothelial and glioma cell migration. *J Biol Chem* 288: 343-54.

KUBOYAMA K, FUJIKAWA A, SUZUKI R, TANGA N, NODA M (2016). Role of Chondroitin Sulfate (CS) Modification in the Regulation of Protein-tyrosine Phosphatase Receptor Type Z (PTPRZ) Activity: PLEIOTROPHIN-PTPRZ-A SIGNALING IS INVOLVED IN OLIGODENDROCYTE DIFFERENTIATION. *J Biol Chem* 29: 18117-18128.

LAMPROPOULOU E, LOGOVITI I, KOUTSIOUMPA M, HATZIAPOSTOLOU M, POLYTARCHOU C, SKANDALIS SS, HELLMAN U, FOUSTERIS M, NIKOLAROPOULOS S, CHOLEVA E, LAMPROU M, SKOURA A, MEGALOOIKONOMOU V, PAPADIMITRIOU E (2018). Cyclin-dependent kinase 5 mediates pleiotrophin-induced endothelial cell migration. *Sci Rep.* 8: 5893.

LAMPROU M, KASPIRIS A, PANAGIOTOPOULOS E, GIANNOUDIS PV, PAPADIMITRIOU E (2014). The role of pleiotrophin in bone repair. *Injury* 45: 1816-1823.

LAMPROU M, KASTANA P, KOFINA F, TZOUPIS H, BARMPOUTSI S, SAJIB MS, KOUTSIOUMPA M, POIMENIDI E, ZOMPRA AA, TASSOPOULOS D, CHOLEVA E, TSELIOS T, MIKELIS CM, PAPADIMITRIOU E (2020). Pleiotrophin selectively binds to vascular endothelial growth factor receptor 2 and inhibits or stimulates cell migration depending on  $\alpha_v\beta_3$  integrin expression. *Angiogenesis* 23: 621-636.

LANDGRAF P, WAHLE P, PAPE H C, GUNDELFINGER E D, KREUTZ M R (2008). The survival-promoting peptide Y-P30 enhances binding of pleiotrophin to syndecan-2 and -3 and supports its neuritogenic activity. *J Biol Chem* 283: 25036–25045.

LEDOUX D, CARUELLE D, SABOURIN J C, LIU J, CREPIN M, BARRITAU D, COURTY J (1997). Cellular distribution of the angiogenic factor heparin affin regulatory peptide (HARP) mRNA and protein in the human mammary gland. *J Histochem Cytochem* 45: 1239–1245.

LEVY JB, CANOLL PD, SILVENNOINEN O, BARNEA G, MORSE B, HONEGGER AM, HUANG JT, CANNIZZARO LA, PARK SH, DRUCK T (1993). The cloning of a receptor-type protein tyrosine phosphatase expressed in the central nervous system. *J Biol Chem* 268: 10573-10581.

LI F, TIAN F, WANG L, WILLIAMSON IK, SHARIFI BG, SHAH PK (2010). Pleiotrophin (PTN) is expressed in vascularized human atherosclerotic plaques: IFN- $\gamma$ /JAK/STAT1 signaling is critical for the expression of PTN in macrophages. *FASEB J* 24: 810-822.

LI YS, GURRIERI M, DEUEL TF (1992). Pleiotrophin gene expression is highly restricted and is regulated by platelet-derived growth factor. *Biochem Biophys Res Commun* 184: 427-432.

MAARTENS AP, BROWN NH (2015). Anchors and signals: the diverse roles of integrins in development. *Curr Top Dev Biol* 112: 233-272.

MARGOLIS RK, RAUCH U, MAUREL P, MARGOLIS RU (1996). Neurocan and phosphacan: two major nervous tissue-specific chondroitin sulfate proteoglycans. *Perspect Dev Neurobiol* 3: 273-90.

MASUDA H, TSUJIMURA A, YOSHIOKA M, ARAI Y, KUBOKI Y, MUKAIT, NAKAMURA T, TSUJI H, NAKAGAWA M, HASHIMOTO-GOTOH T (1997). Bone mass loss due to estrogen

deficiency is compensated in transgenic mice overexpressing human osteoblast stimulating factor-1. *Biochem Biophys Res Commun* 238: 528-533.

McCLAIN C R, SIM F J, GOLDMAN S A (2012). Pleiotrophin suppression of receptor protein tyrosine phosphatase- $\beta/\zeta$  maintains the self-renewal competence of fetal human oligodendrocyte progenitor cells. *J Neurosci Res* 32: 15066–15075.

MIAO J, WANG F, WANG R, ZENG J, ZHENG C, ZHUANG G (2019). Pleiotrophin regulates functional heterogeneity of microglia cells in EAE animal models of multiple sclerosis by activating CCR-7/CD206 molecules and functional cytokines. *Am J Transl Res* 11: 2013-2027.

MICHELOTTI G A, TUCKER A, SWIDERSKA-SYN M, MACHADO M V, CHOI S S, KRUGER L, SODERBLOM E, THOMPSON J W, MAYER-SALMAN M, HIMBURG H A, MOYLAN C A, GUY C D, GARMAN K S, PREMONT R T, CHUTE J P, DIEHL A M (2016). Pleiotrophin regulates the ductular reaction by controlling the migration of cells in liver progenitor niches. *Gut* 65: 683–692.

MIKELIS C, LAMPROU M, KOUTSIOUMPA M, KOUTSIOUBAS A G, SPYRANTI Z, ZOMPRA A A, SPILIOPOULOS N, VRADIS A A, KATSORIS P, SPYROULIAS G A, CORDOPATIS P, COURTY J, PAPADIMITRIOU E (2011). A peptide corresponding to the C-terminal region of pleiotrophin inhibits angiogenesis *in vivo* and *in vitro*. *J Cell Biochem* 112: 1532–1543.

MIKELIS C, SFAELOU E, KOUTSIOUMPA M, KIEFFER N, PAPADIMITRIOU E (2009). Integrin  $\alpha_v\beta_3$  is a pleiotrophin receptor required for pleiotrophin-induced endothelial cell migration through receptor protein tyrosine phosphatase  $\beta/\zeta$ . *FASEB J* 23: 1459–1469.

MILHIET P E, VACHEROT F, CARUELLE J P, BARRITAUULT D, CARUELLE D, COURTY J (1998). Upregulation of the angiogenic factor heparin affin regulatory peptide by progesterone in rat uterus. *J Endocrinol* 158: 389–399.

MITSIADIS T A, SALMIVIRTA M, MURAMATSU T, MURAMATSU H, RAUVALA H, LEHTONEN E, JALKANEN M, THESLEFF I (1995). Expression of the heparin-binding cytokines, midkine (MK) and HB-GAM (pleiotrophin) is associated with epithelial-mesenchymal interactions during fetal development and organogenesis. *Development* 121: 37-51.

MITSIADIS T A, CATON J, DE BARI C, BLUTEAU G (2008). The large functional spectrum of the heparin-binding cytokines MK and HB-GAM in continuously growing organs: The rodent incisor as a model. *Dev Biol* 320: 256–266.

MOURLEVAT S, DEBEIR T, FERRARIO J E, DELBE J, CARUELLE D, LEJEUNE O, DEPIENNE C, COURTY J, RAISMAN-VOZARI R, RUBERG M (2005). Pleiotrophin mediates the neurotrophic effect of cyclic AMP on dopaminergic neurons: Analysis of suppression-subtracted cDNA libraries and confirmation *in vitro*. *Exp Neurol* 194: 243–254.

MUÑOZ-SORIANO V, RUIZ C, PÉREZ-ALONSO M, MLODZIK M, PARICIO N (2013). Nemo regulates cell dynamics and represses the expression of miple, a midkine/pleiotrophin cytokine, during ommatidial rotation. *Dev Biol* 377: 113-125.

MURAMATSU H, ZOU P, KUROSAWA N, ICHIHARA-TANAKA K, MARUYAMA K, INOH K, SAKAI T, CHEN L, SATO M, MURAMATSU T (2006). Female infertility in mice deficient in midkine and pleiotrophin, which form a distinct family of growth factors. *Genes Cells* 11: 1405–1417.

NTENEKOU D, KASTANA P, HATZIAPOSTOLOU M, POLYTARCHOU C, MARAZIOTI A, NIKOU S, HERRADON G, PAPADAKI E, STATHOPOULOS G, MIKELIS CM and PAPADIMITRIOU E (2020). Anaplastic Lymphoma Kinase inhibition as an effective treatment strategy against the enhanced lung carcinogenesis and cancer angiogenesis related to decreased PTPRZ1 expression. *Eur Respir J* 56 (Suppl 64): 3939.

ORR B, VANPOUCKE G, GRACE OC, SMITH L, ANDERSON RA, RIDDICK AC, FRANCO OE, HAYWARD SW, THOMSON AA (2011). Expression of pleiotrophin in the prostate is androgen regulated and it functions as an autocrine regulator of mesenchyme and cancer associated fibroblasts and as a paracrine regulator of epithelia. *Prostate* 71: 305-317.

PALMIERI D, MURA M, MAMBRINI S, PALOMBO D (2015). Effects of Pleiotrophin on endothelial and inflammatory cells: Pro-angiogenic and anti-inflammatory properties and potential role for vascular bio-prosthesis endothelialization. *Adv Med Sci* 60: 287–293.

PANTAZAKA E, PAPADIMITRIOU E (2014). Chondroitin sulfate-cell membrane effectors as regulators of growth factor-mediated vascular and cancer cell migration. *Biochim Biophys Acta Gen Subj* 1840: 2643–2650.

PAPADIMITRIOU E, MIKELIS C, LAMPROPOULOU E, KOUTSIOUMPA M, THEOCHARI K, TSIRMOULA S, THEODOROPOULOU C, LAMPROU M, SFAELOU E, VOURTSIS D BOUDOURIS P (2009). Roles of pleiotrophin in tumor growth and angiogenesis. *Eur Cytokine Netw* 20: 180–190.

PAPADIMITRIOU E, PANTAZAKA E, CASTANA P, TSALIOS T, POLYZOS A, BEIS D (2016). Pleiotrophin and its receptor protein tyrosine phosphatase beta/zeta as regulators of angiogenesis and cancer. *Biochim Biophys Acta Rev Cancer* 1866: 252–265.

PAPADIMITRIOU E, POLYKRATIS A, HATZIAPOSTOLOU M, PARTHYMOU A, POLYTARCHOU C, MIKELIS C (2004). Heparin affin regulatory peptide: a new target for tumour therapy? *Curr Cancer Drug Targets* 4: 471-482.

PAVELIEV M, FENRICH K K, KISLIN M, KUJA-PANULA J, KULESSKIY E, VARJOSALO M, KAJANDER T, MUGANTSEVA E, AHONEN-BISHOPP A, KHIROUG L, KULESSKAYA N, ROUGON G, RAUVALA H (2016). HB-GAM (pleiotrophin) reverses inhibition of neural regeneration by the CNS extracellular matrix. *Sci Rep* 6: 33916.

POIMENIDI E, THEODOROPOULOU C, KOUTSIOUMPA M, SKONDRA L, DROGGITI E, VAN DEN BROEK M, KOOLWIJK P, PAPADIMITRIOU E (2016). Vascular endothelial growth factor A (VEGF-A) decreases expression and secretion of pleiotrophin in a VEGF receptor-independent manner. *Vasc Pharmacol* 80: 11–19.

ROGER J, BRAJEUL V, THOMASSEAU S, HIENOLA A, SAHEL JA, GUILLONNEAU X, GOUREAU O (2006). Involvement of Pleiotrophin in CNTF-mediated differentiation of the late retinal progenitor cells. *Dev Biol* 298: 527-539.

ROSENFELD S M, BOWDEN E T, COHEN-MISSNER S, GIBBY K A, ORY V, HENKE R T, RIEGEL A T, WELLSTEIN A (2012). Pleiotrophin (PTN) Expression and Function and in the Mouse Mammary Gland and Mammary Epithelial Cells. *PLoS One*, 7: e47876.

SAKURAI H, BUSH K T, NIGAM S K (2001). Identification of pleiotrophin as a mesenchymal factor involved in ureteric bud branching morphogenesis. *Development* 128: 3283–3293.

SATO H, FUNAHASHI M, KRISTENSEN D B, TATENO C, YOSHIZATO K (1999). Pleiotrophin as a Swiss 3T3 cell-derived potent mitogen for adult rat hepatocytes. *Exp Cell Res* 246: 152–164.



SHARIFI B G, ZENG Z, WANG L, SONG L, CHEN H, QIN M, SIERRA-HONIGMANN M R, WACHSMANN-HOGIU S, SHAH, P K (2006). Pleiotrophin induces transdifferentiation of monocytes into functional endothelial cells. *Arterioscler Thromb Vasc Biol* 26: 1273–1280.

SHEN D, PODOLNIKOVA NP, YAKUBENKO VP, ARDELL CL, BALABIYEV A, UGAROVA TP, WANG X (2017). Pleiotrophin, a multifunctional cytokine and growth factor, induces leukocyte responses through the integrin Mac-1. *J Biol Chem* 292: 18848-18861.

SONE M, MURAMATSU H, MURAMATSU T, NAKASHIMA T (2011). Morphological observation of the stria vascularis in midkine and pleiotrophin knockout mice. *Auris Nasus Larynx* 38: 41-45.

SVENSSON SL, PASUPULETI M, WALSE B, MALMSTEN M, MÖRGELIN M, SJÖGREN C, OLIN AI, COLLIN M, SCHMIDTCHEN A, PALMER R, EGESTEN A (2010). Midkine and pleiotrophin have bactericidal properties: preserved antibacterial activity in a family of heparin-binding growth factors during evolution. *J Biol Chem* 285: 16105-16115.

TANG C, WANG M, WANG P, WANG L, WU Q, GUO W (2019). Neural Stem Cells Behave as a Functional Niche for the Maturation of Newborn Neurons through the Secretion of PTN. *Neuron* 101: 32–44.

TANGA N, KUBOYAMA K, KISHIMOTO A, KIHARA M, KIYONARI H, WATANABE T, FUJIKAWA A, NODA M (2019). Behavioral and neurological analyses of adult mice carrying null and distinct loss-of-receptor function mutations in protein tyrosine phosphatase receptor type Z (PTPRZ). *PLoS One* 14: e0217880.

TAVELLA S, RUGGIU A, GIULIANI A, BRUN F, CANCIANI B, MANESCU A, MAROZZI K, CILLI M, COSTA D, LIU Y, PICCARDI F, TASSO R, TROMBA G, RUSTICHELLI F,

CANCEDDA R (2012). Bone turnover in wild type and pleiotrophin-transgenic mice housed for three months in the international space station (ISS). *PLoS One* 7: e33179.

THEODORAKOPOULOU O, HATZIAPOSTOLOU M and PAPADIMITRIOU E. ATRA's inhibitory effect on prostate cancer cell growth involves HARP expression (2006). *FEBS J* 273(Suppl 1):86 (PP-35)

VAZIN T, BECKER K G, CHEN J, SPIVAK C E, LUPICA C R, ZHANG Y, WORDEN L, FREED W J (2009). A novel combination of factors, termed SPIE, which promotes dopaminergic neuron differentiation from human embryonic stem cells. *PLoS One* 4: e6606.

WANAKA A, CARROLL S L, MILBRANDT J (1993). Developmentally regulated expression of pleiotrophin, a novel heparin binding growth factor, in the nervous system of the rat. *Dev Brain Res* 72: 133–144.

WANG K, YIN YH, YANG ZQ, YU HF, WANG YS, GUO B, YUE ZP. Hmgb3 Induces the Differentiation of Uterine Stromal Cells Through Targeting Ptn (2019). *Reprod Sci* 26: 891-899.

WANG W, WAN L, CHEN Z, JIN X, LI D (2020). Myofibroblasts control the proliferation of fetal hepatoblasts and their differentiated cholangiocytes during the hepatoblast-to-cholangiocyte transition. *Biochem Biophys Res Commun* 522: 845-851.

WENG T, GAO L, BHASKARAN M, GUO Y, GOU D, NARAYANAPERUMAL J, CHINTAGARI N R, ZHANG K, LIU L (2009). Pleiotrophin regulates lung epithelial cell proliferation and differentiation during fetal lung development via  $\beta$ -catenin and Dlk1. *J Biol Chem* 284: 28021–28032.

WINKLER C, YAO S (2014). The midkine family of growth factors: diverse roles in nervous system formation and maintenance. *Br J Pharmacol* 171: 905-12.

XI G, DEMAMBRO V E, D' COSTA S, XIA S K, COX Z C, ROSEN C J, CLEMMONS D R (2020). Estrogen Stimulation of Pleiotrophin Enhances Osteoblast Differentiation and Maintains Bone Mass in IGFBP-2 Null Mice. *US Endocrinol* 161: 1–13.

XIA Z, OUYANG D, LI Q, LI M, ZOU Q, LI L, YI W, ZHOU E (2019). The Expression, Functions, Interactions and Prognostic Values of PTPRZ1: A Review and Bioinformatic Analysis. *J Cancer* 10: 1663-1674.

YANAGISAWA H, KOMUTA Y, KAWANO H, TOYODA M, SANGO K (2010). Pleiotrophin induces neurite outgrowth and up-regulates growth-associated protein (GAP)-43 mRNA through the ALK/GSK3 $\beta$ / $\beta$ -catenin signaling in developing mouse neurons. *Neurosci Res* 66: 111–116.

YANG S, XUE D DAN, WU B, SUN H MEI, LI X SHUANG, DONG F, LI W SHUAI, JI F QING, ZHOU D SHAN (2013). Pleiotrophin is involved in the amniotic epithelial cell-induced differentiation of human umbilical cord blood-derived mesenchymal stem cells into dopaminergic neuron-like cells. *Neurosci Lett* 539: 86–91.


YU HF, TAO R, YANG ZQ, WANG K, YUE ZP, GUO B (2018). Ptn functions downstream of C/EBP $\beta$  to mediate the effects of cAMP on uterine stromal cell differentiation through targeting Hand2 in response to progesterone. *J Cell Physiol* 233: 1612-1626.

ZHANG L, XIA D, WANG C, GAO F, HU L, LI J, JIN L (2021). Pleiotrophin attenuates the senescence of dental pulp stem cells. *Oral Dis* Online ahead of print.


ZHANG N, YEH H J, ZHONG R, LI Y S, DEUEL T F (1999). A dominant-negative pleiotrophin mutant introduced by homologous recombination leads to germ-cell apoptosis in male mice. *PNAS USA* 96: 6734–6738.

ZHOU Y, FU B, XU X, ZHANG J, TONG X, WANG Y, DONG Z, ZHANG X, SHEN N, ZHAI Y, KONG X, SUN R, TIAN Z, WEI H (2020). PBX1 expression in uterine natural killer cells drives fetal growth. *Sci Transl Med* 12: 1–16.


ZOU P, MURAMATSU H, SONE M, HAYASHI H, NAKASHIMA T, MURAMATSU T (2006). Mice doubly deficient in the midkine and pleiotrophin genes exhibit deficits in the expression of  $\beta$ -tectorin gene and in auditory response. *Lab Invest* 86: 645–653.




**Fig. 1. Timeline representation of PTN discovery.** Important time points of the discovery of the molecule and the different names used during the years. The figure was created with BioRender.com.



**Fig. 2. Schematic representation of the known PTN receptors and their proven biological effects upon PTN binding.** All shown receptors are functional PTN receptors that mainly affect the PTN-mediated regulation of migration of different types of cells. SDC1 and SDC4 have been shown to interact with PTN but there are no data on their involvement in any PTN actions; therefore, they are not presented in this figure. Data on ALK as a PTN receptor are conflicting (see text), but PTN may activate ALK through PTPRZ1 and it seems that ALK mediates some of PTN's actions. NCL conformation and exact mode of localization on the cell surface is not known. The figure was created with BioRender.com.



**Fig. 3. PTPRZ1 interacts with  $\alpha_v\beta_3$  in endothelial cells independently of PTN.** Serum-starved human umbilical vein endothelial cells were incubated with PTN (100 ng/ml) for 10 min. Formation of  $\beta_3$ -PTPRZ1 complexes as evidenced by *in situ* proximity ligation assay are shown on the left. The box plots shown on the right indicate the median, mean and range of the detected signals (n = 10-12 image fields with 4-5 cells per image per sample type, each sample run in duplicate). Details on the methodology used can be found at Koutsoumpa *et al.*, 2015. Scale bar corresponds to 10  $\mu$ m.



**Fig. 4.** Schematic representation of the up to date known PTN-induced pathway in endothelial cells. PTN has a moderate stimulatory effect on endothelial cell migration via PTPRZ1/ $\alpha_v\beta_3$ /cell surface NCL (Mikelis *et al.*, 2009; Koutsoumpa *et al.*, 2013) but an inhibitory effect through VEGFR2 (Lamprou *et al.*, 2020). The figure was created with BioRender.com.